

Long-Term Regulation of Renal Urea Transporters during Antidiuresis

Dong-Un Kim, M.D.

Department of Pediatrics, College of Medicine, The Catholic University of Korea, Seoul, Korea

To produce a concentrated urine, the renal medulla needs hypertonicity for the reabsorption of free water from collecting duct. The single effect that increases interstitial tonicity in the outer medulla is the active NaCl reabsorption in the thick ascending limb, while the single effect in the inner medulla is the passive efflux of NaCl through the thin ascending limb. The passive mechanism in the inner medulla requires high interstitial urea concentration. Two main groups of urea transporters (UT-A, UT-B) are present in the kidney, which maintains the high concentration of urea in the deepest portion of the inner medulla by intra-renal urea recycling. Recent studies suggest that UT-A1 in the terminal inner medullary collecting duct is up-regulated when urine or inner medullary interstitial urea is depleted in order to enhance the reabsorption of urea, while UT-A2 in the descending thin limb of loops of Henle and UT-B in the descending vasa recta are increased when outer medullary interstitial urea concentration is high, in order to prevent the loss of urea from the medulla to the systemic circulation, thereby increasing intra-renal urea recycling. This review will summarize the functions of the renal urea transporters in urine concentration mechanism and the recent knowledge about their long-term regulation.

Key Words : Urine concentration, Water, Urea, Urea transporters

The urea transporter family

The urea transporter family includes two main groups: UT-A and UT-B¹⁾. The four isoforms of rat UT-A (UT-A1-UT-A4) are found in the kidney, which are produced by alternate splicing of mRNA from a single gene²⁻⁴⁾. UT-A1 is the longest isoform and the others are composed of part of UT-A1. UT-A2 is basically the C-terminal half of UT-A1, while UT-A3 is basically the N-terminal half of UT-A1^{4, 5)}. UT-A4 basically consists of the N-terminal quarter of UT-A1 spliced to the C-terminal quarter of UT-A1⁴⁾. UT-A1 and UT-A3 are localized in the IMCD^{6, 8)}, and UT-A2 is located in the descending thin limb of

Henle's loop^{9, 10)}. The precise tubular location of UT-A4 is unknown.

UT-B is encoded by a different gene from that encoding UT-A1 and was first cloned from a human bone marrow library¹¹⁾ and subsequently isolated from a rat inner medullary library by homology screening^{12, 13)}. UT-B is expressed in the endothelial cells of the descending vasa recta and the membrane of red blood cells¹¹⁻¹³⁾.

Urine concentration

While NaCl is actively reabsorbed from the thick ascending limb in the outer medulla, there is no active process of NaCl reabsorption in the inner medulla. However, NaCl is reabsorbed from the thin ascending limb by virtue of the presence of urea in the inner medullary interstitium. The passive countercurrent multiplication explains the urine concentrating mecha-

Corresponding author: Dong-Un Kim, M.D.
Department of Pediatrics, College of Medicine, The Catholic University of Korea, Seoul, Korea
Tel: 031)820-3580, Fax: 031)821-3108
E-mail: dukim@catholic.ac.kr

nism in mammals by which inner medullary tonicity is increased without active transport^{14, 15}. In this theory, the higher urea concentration within the inner medullary interstitial fluid osmotically balances the higher NaCl concentration in the lumen of the thin ascending limb. Since the thin ascending limb is impermeable to water, highly permeable to NaCl, but only somewhat permeable to urea, NaCl is passively reabsorbed from the lumen of thin ascending limb to inner medullary interstitium by electrochemical gradient (Fig. 1).

Intra-renal urea recycling

The source of urea in the inner medullary interstitial fluid is reabsorption from the terminal inner medullary collecting duct. UT-A1 urea transporter is present in the terminal inner medullary collecting duct cells and facilitates the delivery of urea from collecting duct lumen to inner medullary interstitium. However, the highly concentrated urea in the terminal inner medulla could not be achieved if urea was rapidly taken up by the fenestrated ascending vasa recta and returned to

the systemic circulation. UT-A2 in the descending thin limb of loops of Henle and UT-B in descending vasa recta enable a large fraction of the urea carried in the ascending blood to be reintroduced into the deep inner medulla. This creates an intra-renal urea recycling process that minimizes the loss of urea from the renal medulla and maintains the medullary osmotic gradient (Fig. 1).

Long term regulation of urea transporters

1. UT-A1

In Diabetes mellitus (DM), unabsorbed glucose retains water in the lumen and causes severe osmotic diuresis. In rats with uncontrolled DM (induced by streptozotocin), UT-A1 urea transporter and AQP2 water channel abundances in the collecting duct were significantly increased after 10-21 days after streptozotocin injection¹⁶⁻¹⁸, suggesting that these transporter proteins are up-regulated during DM-induced osmotic diuresis in order to prevent a serious volume depletion.

On the other hand, vasopressin rapidly increases the phosphorylation of UT-A1¹⁹ and is known to be elevated in DM^{20, 21}, which bring the thought that the increase of UT-A1 abundance in diabetic rats may be due to their high plasma vasopressin level.

To explore the effect of vasopressin on UT-A1 abundance in diabetic rats, Brattleboro rats (which lack circulating vasopressin) were infused with vasopressin or made diabetic by streptozotocin injection or received both procedures together²². DM for 10 days did not increase UT-A1 abundance in Brattleboro rats devoid of vasopressin, while vasopressin infusion (for 12 days) did significantly increase UT-A1 abundance, compared to untreated Brattleboro rats. DM combined with vasopressin infusion had more increase in UT-A1 abundance over vasopressin infusion alone. These results suggest that vasopressin is necessary for the increase of UT-A1 abundance during diabetes. It is also suggested that a factor other than vasopressin is responsible for the increase in UT-A1 protein abun-




Fig. 1. The location of urea transporters and the intra-renal urea recycling. UT, urea transporters; AQP, aquaporin (modified from Ref. 26 with permission). NaCl is actively transported out from thick ascending limb in the outer medulla, while passively transported out from thin ascending limb in the inner medulla. The passive transport of NaCl in the inner medulla requires high concentration of urea in the inner medullary interstitium, which is maintained by urea transporters that mediate intra-renal recycling of urea. The efflux of NaCl from both thick and thin ascending limb increases medullary interstitial tonicity and then water is reabsorbed from collecting duct through AQP 2-4 water channels.

dance because the same amount of vasopressin was administered to the two vasopressin treated groups of Brattleboro rats²².

Urea is the major urinary solute in mammals, comprising 40–50% of total urinary solute in rats on a regular diet (containing 1% NaCl and 23% protein). Rats with streptozotocin-induced DM have an increase in absolute urea excretion, but the relative amount (percent) of urea in total urinary solute is actually decreased (to below 30%) due to the marked increase in glucose excretion¹⁷. The ongoing osmotic diuresis due to non-reabsorbable glucose causes water to be retained in the tubule lumen which dilutes the urine urea concentration. Several urea-specific signaling pathways have been identified in cultured mIMCD3 cells and the renal medulla^{23, 24}, suggesting the possibility that changes in the percentage or concentration of urea in the urine or medullary interstitium could be a factor that regulates UT-A1 protein abundance.

To prove our hypothesis, diabetic rats were fed 10% urea diet in order to increase urea excretion, thereby keeping the percent of urea in total urinary solute at the control level. The urine urea concentration in this group was significantly increased when compared to that of diabetic rats with regular diet, although it was reduced when compared to control. UT-A1 abundance did not change in urea fed diabetic rats, which suggests that the increase of UT-A1 in diabetic rats is due to the decrease of urea in the urine²⁵.

To further examine our hypothesis, rats were fed high salt diet containing 4% NaCl to induce NaCl diuresis. An additional group of rats were fed 4% NaCl and 3% urea in order to keep the percent of urea in total urinary solute at the control level. The absolute amount of urea excretion in the rats with 4% NaCl diet was not different from that of control rats, but the percent of urea in total urinary solute and urine urea concentration were significantly decreased due to the markedly increased NaCl excretion. UT-A1 abundance in the rats with 4% NaCl diet was significantly increased but not in the rats with 4% NaCl

plus 3% urea diet²⁵.

Next, we fed rats 5% urea diet in order to increase the percent of urea in total urinary solute (to over 60%) by increasing urea excretion. Usually, low urine urea concentrations and low urine osmolalities occur simultaneously, making it difficult to know which one is responsible for an increase in UT-A1 abundance. However, these two conditions can be dissociated if total urinary solute is increased by urea itself. In the rats with 5% urea diet, the urine urea concentration was increased, but the urine osmolality was decreased, which suggests that too much urea is not helpful for urine concentrating ability; rather it inhibits urine concentration by acting as an osmotic agent. Urea diuresis did not change UT-A1 abundance despite the decrease in urine osmolality and the presence of osmotic diuresis, suggesting that UT-A1 increases in response to a reduced percentage of urea in urine and/or a reduced urine urea concentration, rather than to a low urine osmolality²⁵.

2. UT-A2 and UT-B

UT-A2 and UT-B mediate intra-renal urea recycling through the thin descending limb and descending vasa recta, respectively^{26, 27}. UT-A2 is up-regulated in UT-B knock-out mice, suggesting that loss of one urea recycling pathway can be partially compensated for by increasing the other pathway²⁸. Likewise, UT-B mRNA is up-regulated in UT-A2 knock-out mice, even though the increase was not significant²⁹. UT-A2 protein or mRNA has been increased in dehydrated Sprague-Dawley rats^{3, 30} or DDAVP-infused Brattleboro rats^{9, 10}, conditions where urine urea concentration is increased. UT-A2 and UT-B abundances were increased in urea diuresis²⁵, where the urine urea concentration increased, but was unchanged or decreased in NaCl diuresis and DM-induced glucose diuresis^{18, 25}, where the urine urea concentration is low. This suggests that UT-A2 and UT-B abundances increase when the medullary interstitial urea concentration is high, which would tend to increase

intra-renal urea recycling during antidiuresis, thereby preventing the loss of urea from the medulla and maintaining medullary interstitial osmolality²⁵⁾.

References

- 1) Sands JM, Timmer RT, Gunn RB : Urea transporters in kidney and erythrocytes. *Am J Physiol Renal Physiol* **273**:F321-F339, 1997
- 2) You G, Smith CP, Kanai Y, Lee WS, Stelzner M, Hediger MA : Cloning and characterization of the vasopressin-regulated urea transporter. *Nature* **365**: 844-847, 1993
- 3) Smith CP, Lee WS, Martial S, Knepper MA, You G, Sands JM, Hediger MA : Cloning and regulation of expression of the rat kidney urea transporter (rUT2). *J Clin Invest* **96**:1556-1563, 1995
- 4) Karakashian A, Timmer RT, Klein JD, Gunn RB, Sands JM, Bagnasco SM : Cloning and characterization of two new isoforms of the rat kidney urea transporter: UT-A3 and UT-A4. *J Am Soc Nephrol* **10**:230-237, 1999
- 5) Shayakul C, Tsukaguchi H, Berger UV, Hediger MA : Molecular characterization of a novel urea transporter from kidney inner medullary collecting ducts. *Am J Physiol Renal Physiol* **280**:F487-F494, 2001
- 6) Nielsen S, Terris J, Smith CP, Hediger MA, Ecelbarger CA, Knepper MA : Cellular and subcellular localization of the vasopressin-regulated urea transporter in rat kidney. *Proc Natl Acad Sci USA* **93**:5495-5500, 1996
- 7) Shayakul C, Knepper MA, Smith CP, DiGiovanni SR, and Hediger MA : Segmental localization of urea transporter mRNAs in rat kidney. *Am J Physiol Renal Physiol* **272**:F654-F660, 1997
- 8) Terris J, Knepper MA, Wade JB : UT-A3 : localization and characterization of an additional urea transporter isoform in the IMCD. *Am J Physiol Renal Physiol* **280**:F325-F332, 2001
- 9) Shayakul C, Smith CP, Mackenzie HS, Lee WS, Brown D, Hediger MA : Long-term regulation of urea transporter expression by vasopressin in Brattleboro rats. *Am J Physiol Renal Physiol* **278**: F620-F627, 2000
- 10) Wade JB, Lee AJ, Liu J, Ecelbarger CA, Mitchell C, Bradford AD, Terris J, Kim GH, Knepper MA : UT-A2 : a 55-kDa urea transporter in thin descending limb whose abundance is regulated by vasopressin. *Am J Physiol Renal Physiol* **278**:F52-F62, 2000
- 11) Olives B, Neau P, Bailly P, Hediger MA, Rousset G, Carton JP, Ripoché P : Cloning and functional expression of a urea transporter from human bone marrow cells. *J Biol Chem* **269**:31649-31652, 1994
- 12) Tsukaguchi H, Shayakul C, Berger UV, Tokui T, Brown D, Hediger MA : Cloning and characterization of the urea transporter UT3 : localization in rat kidney and testis. *J Clin Invest* **99**:1506-1515, 1997
- 13) Xu Y, Olives B, Bailly P, Fischer E, Ripoché P, Ronco P, Carton JP, Rondeau E : Endothelial cells of the kidney vasa recta express the urea transporter HUT11. *Kidney Int* **51**:138-146, 1997
- 14) Kokko JP, Rector FC Jr : Countercurrent multiplication system without active transport in inner medulla. *Kidney Int* **2**:214-223, 1972
- 15) Stephenson JL : Concentration of urine in a central core model of the renal counterflow system. *Kidney Int* **2**:85-94, 1972
- 16) Nejsum LN, Kwon TH, Marples D, Flyvbjerg A, Knepper MA, Frokiaer J, Nielsen S : Compensatory increase in AQP2, p-AQP2, and AQP3 expression in rats with diabetes mellitus. *Am J Physiol Renal Physiol* **280**:F715-F726, 2001
- 17) Kim DU, Sands JM, and Klein JD : Changes in renal medullary transport proteins during uncontrolled diabetes mellitus in rats. *Am J Physiol Renal Physiol* **285**:F303-F309, 2003
- 18) Bardoux P, Ahloulay M, Le Maout S, Bankir L, Trinh-Trang-Tan MM : Aquaporin-2 and urea transporter-A1 are upregulated in rats with type I diabetes mellitus. *Diabetologia* **44**:637-645, 2001
- 19) Zhang C, Sands JM, and Klein JD : Vasopressin rapidly increases the phosphorylation of the UT-A1 urea transporter activity in rat IMCDs through PKA. *Am J Physiol Renal Physiol* **282**:F85-F90, 2002
- 20) Brooks DD, Nutting DF, Crofton JT, Share L : Vasopressin in rats with genetic and streptozotocin-induced diabetes. *Diabetes* **38**:54-57, 1989
- 21) Trinder D, Phillips PA, Stephenson JM, Risvanis J, Aminian A, Adam W, Cooper M, Johnston CI : Vasopressin V1 and V2 receptors in diabetes mellitus. *Am J Physiol Endocrinol Metab* **266**:E217-E223, 1994
- 22) Kim D-U, Sands JM, Klein JD : Role of vasopressin in diabetes mellitus-induced changes in medullary transport proteins involved in urine concentration in Brattleboro rats. *Am J Physiol Renal Physiol* **286**: F760-F766, 2004
- 23) Berl T : On the adaptation of renal cells to hypertonicity. *Am J Kidney Dis* **35**:XLVII-XLVII, 2000
- 24) Tian W, Cohen DM : Signaling and gene regulation by urea in cells of the mammalian kidney medulla. *Comp Biochem Physiol A* **130**:429-436, 2001
- 25) Kim D-U, Klein JD, Racine S, Murrell BP, Sands JM : Urea may regulate urea transporter protein abundance during osmotic diuresis. *Am J Physiol Renal Physiol* **288**:F188-F197, 2005
- 26) Sands JM : Molecular approaches to urea trans-

- porters. *J Am Soc Nephrol* **13**:2795-2806, 2002
- 27) Sands JM : Molecular mechanisms of urea transport. *J Membr Biol* **191**:149-163, 2003
- 28) Klein JD, Sands JM, Qian L, Wang X, Yang B : Upregulation of urea transporter UT-A2 and water channels AQP2 and AQP3 in mice lacking urea transporter UT-B. *J Am Soc Nephrol* **15**:1161-1167, 2004
- 29) Uchida S, Sohara E, Rai T, Ikawa M, Okabe M, Sasaki S : Impaired urea accumulation in the inner medulla of mice lacking the urea transporter UT-A2. *Mol Cell Biol* **25**:7357-7363, 2005
- 30) Bagnasco SM, Peng T, Nakayama Y, Sands JM : Differential expression of individual UT-A urea transporter isoforms in rat kidney. *J Am Soc Nephrol* **11**:1980-1986, 2000