

Hypernatremia : Successful Treatment

Soo Wan Kim, M.D.

Department of Internal Medicine, Chonnam National University Medical School, Gwangju, Korea

Hypernatremia reflects a net water loss or a hypertonic sodium gain, with inevitable hyperosmolality. Severe symptoms are usually evident only with acute and large increases in plasma sodium concentrations to above 158-160 mmol/l. Importantly, the sensation of intense thirst that protects against severe hypernatremia in health may be absent or reduced in patients with altered mental status or with hypothalamic lesions affecting their sense of thirst and in infants and elderly people. Non-specific symptoms such as anorexia, muscle weakness, restlessness, nausea, and vomiting tend to occur early. More serious signs follow, with altered mental status, lethargy, irritability, stupor, and coma. Acute brain shrinkage can induce vascular rupture, with cerebral bleeding and subarachnoid hemorrhage. However, in the vast majority of cases, the onset of hypertonicity is low enough to allow the brain to adapt and thereby to minimize cerebral dehydration. Organic osmolytes accumulated during the adaptation to hypernatremia are slow to leave the cell during rehydration. Therefore, if the hypernatremia is corrected too rapidly, cerebral edema results as the relatively more hypertonic ICF accumulates water. To be safe, the rate of correction should not exceed 12 mEq/liter/day.

Key Words : Hypernatremia, Cell volume regulation, Diabetes insipidus

Hypernatremia, defined as a rise in the serum sodium concentration to a value exceeding 145 mmol per liter, is a common electrolyte disorder. Because sodium is a functionally impermeable solute, it contributes to tonicity and induces the movement of water across cell membranes. Therefore, hypernatremia invariably denotes hypertonic hyperosmolality and always causes cellular dehydration, at least transiently. Although correction of transient increases in plasma osmolality is usually well tolerated, correction of chronic plasma hypertonicity with rehydration therapy may be accompanied by brain swelling, herniation, and death¹. The clinical differences between acute and chronic osmolar disorders can be understood through a consideration of the different mechanisms by which cells in the brain regulate their volume in response to brief and sustained osmotic challenges².

Pathophysiology

As in other types of cells, hypertonically stressed brain cells regulate their volume initially by the rapid uptake of electrolytes³. With a prolonged elevation of plasma osmolality, however, most excess electrolytes in the brain are replaced by organic solutes. These solutes have historically been termed "idiogenic osmoles" because they were unidentified and thought to be produced by the brain cells themselves⁴. Studies in animals and cultured cells⁵ have revealed that idiogenic osmoles are the same organic osmolytes used by all organisms for volume regulation. The most important organic osmolytes in the mammalian brain include myo-inositol, taurine, glycerylphosphorylcholine, and betaine, which are accumulated primarily by uptake from extracellular fluids through the activation of sodium-dependent cotransporters⁵.

Beginning on the first day of the hypernatremia, brain volume is largely restored due noth to water

Correspondence author : Soo Wan Kim, M.D.
Department of Internal Medicine, Chonnam National University
Medical School, Gwangju, Korea
Tel : 062)220-6272, Fax : 062)220-8578
Email : skimw@chonnam.ac.kr

movement from the cerebrospinal fluid into the brain (thereby increasing the interstitial volume)⁵⁾ and to the uptake of solutes by the cells (thereby pulling water into the cells and restoring the cell volume)⁵⁻⁷⁾. The latter response involves an initial uptake of sodium and potassium salts, followed by the later accumulation of osmolytes, which in animals consists primarily of myo-inositol and the amino acids glutamine and glutamate^{6,7)}. Myo-inositol is taken up from the extracellular fluid via an increase in the number of sodium-myoinositol cotransporters in the cell membrane⁸⁾, whereas the source (uptake from the extracellular fluid or production within the cells) of glutamine and glutamate is at present unknown.

The cerebral adaptation in hyponatremia has two important clinical consequences: Chronic hyponatremia is much less likely to induce neurologic symptoms. Assessment of symptoms attributable to hyponatremia is often difficult because most affected adults have underlying neurologic disease. The latter is required to diminish the protective thirst mechanism that normally prevents the development of hyponatremia, even in patients with diabetes insipidus. Correction of chronic hyponatremia must occur slowly to prevent rapid fluid movement into the brain and cerebral edema, changes that can lead to seizures and coma⁹⁾. Although the brain cells can rapidly lose potassium and sodium in response to this cell swelling, the loss of accumulated osmolytes occurs more slowly, a phenomenon that acts to hold water within the cells⁵⁾. The loss of myoinositol, for example, requires both a reduction in synthesis of new sodium-inositol cotransporters⁸⁾ and the activation of a specific inositol efflux mechanism in the cell membrane¹⁰⁾. The delayed clearance of osmolytes from the cell can predispose to cerebral edema if the plasma sodium concentration is lowered too rapidly. As a result, the rate of correction in asymptomatic patients should not exceed 12 mEq/L per day, which represents an average of 0.5 mEq/L per hour.

Causes of hyponatremia

Hyponatremia represents a deficit of water in relation to the body's sodium stores, which can result from a net water loss or a hypertonic sodium gain (Table 1). Net water loss accounts for the majority of cases of hyponatremia¹⁾. It can occur in the absence of a sodium deficit (pure water loss) or in its presence (hypotonic fluid loss). Hypertonic sodium gain usually results from clinical interventions or accidental sodium loading. Because sustained hyponatremia can occur only when thirst or access to water is impaired, the groups at highest risk are patients with altered mental status, intubated patients, infants, and elderly persons¹¹⁾. Hyponatremia in infants usually results from diarrhea, whereas in elderly persons it is usually associated with infirmity or febrile illness^{12,13)}. Thirst impairment also occurs in elderly patients¹⁴⁾. Often the cause is evident from the history. Measurement of

Table 1. Major Causes of Hyponatremia

Euvolemic Hyponatremia (Pure water loss)
Extrarenal losses
Respiratory (tachycardia)
Dermal (sweating, fever)
Renal losses
Central diabetes insipidus
Nephrogenic diabetes insipidus
Other
Inability to gain access to fluids
Hypodipsia or adipsia
Reset osmostat (essential hyponatremia)
Hypovolemic Hyponatremia (water Deficit in Excess of Sodium Deficit)
Extrarenal losses
Gastrointestinal losses (e.g., diarrhea, vomiting, fistulas)
Dermal (burns, excessive sweating)
Renal losses
Osmotic diuresis (mannitol, glucose, urea)
Loop diuretics
Postobstructive diuresis
Intrinsic renal disease
Hypervolemic Hyponatremia (Sodium gain in Excess of Water Gain)
Hypertonic saline or NaHCO ₃ administration
Infants or comatose patients given hypertonic feedings
Mineralocorticoid excess

urine osmolality in relation to the plasma osmolality and the urine sodium concentration help if the cause is unclear (Fig. 1). Patients with diabetes insipidus present with polyuria and polydipsia (and not hypernatremia unless thirst sensation is impaired). Central diabetes insipidus and nephrogenic diabetes insipidus may be differentiated by the response to water deprivation (failure to concentrate urine) followed by the V₂-receptor agonist desmopressin, causing concentration of urine in patients with central diabetes insipidus.

Clinical manifestations

Signs and symptoms of hypernatremia largely reflect central nervous system dysfunction and are prominent when the increase in the serum sodium concentration is large or occurs rapidly (i.e., over a period of hours)¹⁵. Most outpatients with hypernatremia are either very young or very old¹⁶. Common symptoms in infants include hyperpnea, muscle weakness, restlessness, a characteristic high-pitched cry, insomnia, lethargy, and even coma. Convulsions are typically absent except in cases of inadvertent sodium loading or aggressive rehydration^{17, 18}. Unlike infants, elderly patients generally have few symptoms until the serum sodium concentration exceeds 160 mmol per liter¹⁹. Intense thirst may be present initially, but it

dissipates as the disorder progresses and is absent in patients with hypodipsia. The level of consciousness is correlated with the severity of the hypernatremia²⁰. Muscle weakness, confusion, and coma are sometimes manifestations of coexisting disorders rather than of the hypernatremia itself (Table 2).

Brain shrinkage induced by hypernatremia can cause vascular rupture, with cerebral bleeding, sub-arachnoid hemorrhage, and permanent neurologic damage or death. Brain shrinkage is countered by an adaptive response that is initiated promptly and consists of solute gain by the brain that tends to restore lost water. This response leads to the normalization of brain volume and accounts for the milder symptoms of hypernatremia that develops slowly²¹. However, the normalization of brain volume does not correct hyperosmolality in the brain. In patients with prolonged

Table 2. Signs and Symptoms of Hypermnatemia

Depression of sensorium
Irritability
Seizures (unusual in adults)
Focal neurologic deficits
Muscle spasticity (unusual in adults)
Signs of volume depletion (variable)
Fever
Nausea or vomiting
Labored respiration
Intense thirst




Fig. 1. Diagnostic algorithm for hypernatremia.

hyperosmolality, aggressive treatment with hypotonic fluids may cause cerebral edema, which can lead to coma, convulsions, and death^{17, 18)}.

Treatments

The primary goal in the treatment of patients with hyponatremia is the restoration of serum tonicity. In patients with hyponatremia that has developed over a period of hours, rapid correction of plasma sodium (falling by 1 mmol/L per hour) improves the prognosis without the risk of convulsions and cerebral edema¹⁾. Management of a shocked patient needs specialist input and close monitoring, preferably in a high dependency unit. Intravenous normal saline should be used to correct the extracellular fluid depletion, with calculation of the free water deficit to determine how much 5% dextrose to give. In patients with hyponatremia of longer or unknown duration, reducing the sodium concentration more slowly is prudent. Patients should be given intravenous 5% dextrose for acute hyponatremia or half-normal saline (0.45% sodium chloride) for chronic hyponatremia if unable to tolerate oral water. Central diabetes insipidus is treated with desmopressin, either as intranasal spray or tablets, with careful monitoring to avoid the complications of water intoxication (delaying one dose each week to allow polyuria and thirst to “break-through” in patients susceptible to hyponatremia with desmopressin may be prudent). Treatment of nephrogenic diabetes insipidus includes removal of precipitating drugs (if possible) and sometimes initiation of thiazide diuretics, non-steroidal anti-inflammatory drugs, or both. The following discussion primarily applies to the majority of patients in whom hyponatremia is induced by water loss.

1. Water deficit

The water deficit in the hyponatremic patient can be estimated from the following formula^{22, 23)}.

$$\text{Water deficit} = \text{CBW} \times \left(\frac{\text{plasma } [\text{Na}^+]}{140} - 1 \right)$$

CBW refers to estimated current body water. The total body water is normally about 60 and 50 percent of lean body weight in younger men and women, respectively, and is somewhat lower in the elderly (about 50 and 45 percent in men and women, respectively)²³⁾. However, it is probably reasonable to use values about 10 percent lower (50 and 40 percent) in hyponatremic patients who are water-depleted. Thus, in a 60 kg woman with a plasma sodium concentration of 168 mEq/L, total body water is about 40 percent of body weight and the water deficit can be approximated from:

$$\text{Water deficit} = 0.4 \times 60 \left(\frac{168}{140} - 1 \right) = 4.8 \text{ liters}$$

This formula estimates the amount of positive water balance required to return the plasma sodium concentration to 140 mEq/L. Then, when calculating the amount of free water to give (either intravenously, as dextrose in water, or orally if the patient is able to drink), insensible losses and some part of urine and gastrointestinal losses must be added to the calculation. Although this formula provides an adequate estimate of the water deficit in patients with hyponatremia caused by pure water loss, it underestimates the deficit in patients with hypotonic fluid loss.

2. Rate of correction

There are no definitive clinical trials, but data in children suggest that the maximum safe rate at which the plasma sodium concentration should be lowered is by 0.5 mEq/L per hour and no more than by 12 mEq/L per day²⁴⁾. Overly rapid correction is potentially dangerous in hyponatremia (as it is in hyponatremia). Hyponatremia initially causes fluid movement out of the brain and cerebral contraction that is primarily responsible for the associated symptoms. Within one to three days, however, brain volume is largely restored due to water movement from the cerebrospinal fluid into the brain and to the uptake of solutes by the

cells²⁴). Rapidly lowering the plasma sodium concentration once this adaptation has occurred causes osmotic water movement into the brain, increasing brain size above normal. This cerebral edema can then lead to seizures, permanent neurologic damage, or death^{17, 18}). This sequence of an adverse response to therapy has been primarily described in children in whom the hyponatremia was corrected at a rate exceeding 0.7 mEq/L per h²²). In comparison, no neurologic sequelae were induced if the plasma sodium concentration were lowered at 0.5 mEq/L per hour²³).

3. Common errors in management

Isotonic saline is unsuitable for correcting hyponatremia. Consider a 50-year-old man with a serum sodium concentration of 162 mmol per liter and a body weight of 70 kg (estimated volume of total body water, 42 liters [0.5×70]). The retention of 1 liter of 0.9 percent sodium chloride will decrease the serum sodium concentration by only 0.22 mmol per liter ($[(154-162) \div (35+1)] = -0.22$). Although the sodium concentration of the infusate is lower than the patient's serum sodium concentration, it is not sufficiently low to alter the hyponatremia substantially. Furthermore, ongoing hypotonic fluid losses might outpace the administration of isotonic saline, aggravating the hyponatremia. The sole indication for administering isotonic saline to a patient with hyponatremia is a depletion of extracellular-fluid volume that is sufficient to cause substantial hemodynamic compromise. Even in this case, after a limited amount of isotonic saline has been administered to stabilize the patient's circulatory status, a hypotonic fluid (i.e., 0.2 percent or 0.45 percent sodium chloride) should be substituted in order to restore normal hemodynamic values while correcting the hyponatremia. If a hypotonic fluid is not substituted for isotonic saline, the extracellular-fluid volume may become seriously overloaded.

Extreme care must be taken to avoid excessively rapid correction or overcorrection of hyponatremia, which increases the risk of iatrogenic cerebral edema,

with possibly catastrophic consequences. Selecting the most hypotonic infusate that is suitable for the particular type of hyponatremia ensures the administration of the least amount of fluid. Appropriate allowances for ongoing fluid losses must be made to prevent serious deviations in either direction from the targeted serum sodium concentration. Scrupulous adherence to these management guidelines should help prevent such complications. Most importantly, the fluid prescription should be reassessed at regular intervals in the light of laboratory values and the patient's clinical status¹).

Prognosis

The morbidity and mortality of patients with acute hyponatremia is very high. In children, the mortality of acute hyponatremia ranges between 10 and 70%. As many as two-thirds of the survivors have neurologic sequelae. In contrast, mortality in chronic hyponatremia is 10%. In adults, serum Na concentrations above 160 mmol/L are associated with a 75% mortality¹). Since hyponatremia in adults occurs in the setting of serious disease the mortality rates may reflect the severity of the underlying diseases rather than that of hyponatremia.

References

- 1) Adrogue HJ, Madias NE: Hyponatremia. *New Engl J Med* **342**:1493-1499, 2000
- 2) Petis LD, Luchitti A, Emma F: Cell volume regulation and transport mechanisms across the blood-brain barrier: implications for the management of hyponatremic states. *Eur J Pediatrics* **160**:71-77, 2001
- 3) McManus ML, Strange K: Acute volume regulation of brain cells in response to hypertonic challenge. *Anesthesiology* **78**:1132-1137, 1993
- 4) McDowell ME, Wolf AV, Steer A: Osmotic volumes of distribution: idiogenic changes in osmotic pressure associated with administration of hypertonic solutions. *Am J Physiol* **180**:545-558, 1955
- 5) Strange K: Regulation of solute and water balance and cell volume in the central nervous system. *J Am*

- Soc Nephrol* **3**:12-27, 1992
- 6) Heilig CW, Stromski ME, Blumenfeld JB, Lee JP, Gullan SR: Characterization of the major brain osmolytes that accumulate in salt-loaded rats. *Am J Physiol* **257**:F1108-F1116, 1989
 - 7) Lien YH, Shapiro JI, Chan L: Effect of hypernatremia on organic brain osmoles. *J Clin Invest* **85**:1427-1435, 1990
 - 8) Paredes A, McManus M, Kwon HM, Strange K: Osmoregulation of Na⁺-inositol cotransporter activity and mRNA levels in brain glial cells. *Am J Physiol* **263**:C1282-C1288, 1992
 - 9) Lee JH, Arcinue E, Ross BD: Brief report: Organic osmolytes in the brain of an infant with hypernatremia. *N Engl J Med* **331**:439-442, 1994
 - 10) Strange K, Morrison R, Shrode L, Putnam R: Mechanism and regulation of swelling-activated inositol efflux in brain glial cells. *Am J Physiol* **265**:C244-C256, 1993
 - 11) Gennari FJ: Hypo-hypernatraemia: disorders of water balance. In: Oxford textbook of clinical nephrology. 2nd ed., edited by Davison AM, Cameron JS, Gennari FJ, Kerr DNS, Ritz E, Winearls CG, New York, Oxford University Press, 1998, p175-200.
 - 12) Finberg L, Harrison HE: Hyponatremia in infants: an evaluation of the clinical and biochemical findings accompanying this state. *Pediatrics* **16**:1-14, 1955
 - 13) Bruck E, Abal G, Aceto T Jr: Pathogenesis and pathophysiology of hypertonic dehydration with diarrhea: a clinical study of 59 infants with observations of respiratory and renal water metabolism. *Am J Dis Child* **115**:122-144, 1968
 - 14) Phillips PA, Bretherton M, Johnston CI, Gray L: Reduced osmotic thirst in healthy elderly men. *Am J Physiol* **261**:R166-R171, 1991
 - 15) Snyder NA, Feigal DW, Arief AI: Hyponatremia in elderly patients: a heterogeneous, morbid, and iatrogenic entity. *Ann Intern Med* **107**:309-319, 1987
 - 16) Arief AI, Guisado R: Effects on the central nervous system of hypernatremic and hyponatremic states. *Kidney Int* **10**:104-116, 1976
 - 17) Morris-Jones PH, Houston IB, Evans RC: Prognosis of the neurological complications of acute hypernatraemia. *Lancet* **2**:1385-1389, 1967
 - 18) Hogan GR, Dodge PR, Gill SR, Master S, Sotos JF: Pathogenesis of seizures occurring during restoration of plasma tonicity to normal in animals previously chronically hypernatremic. *Pediatrics* **43**:54-64, 1969
 - 19) Hiromatsu K, Kobayashi T, Fujii N, Itoyama Y, Goto I, Murakami J: Hyponatremic myopathy. *J Neurol Sci* **122**:144-147, 1994
 - 20) De Villota ED, Cavanilles JM, Stein L, Shubin H, Weil MH: Hyperosmolar crisis following infusion of hypertonic sodium chloride for purposes of therapeutic abortion. *Am J Med* **55**:116-122, 1973
 - 21) Lien YH, Shapiro JI, Chan L: Effects of hypernatremia on organic brain osmoles. *J Clin Invest* **85**:1427-1435, 1990
 - 22) Kahn A, Brachet E, Blum D: Controlled fall in natremia and risk of seizures in hypertonic dehydration. *Intensive Care Med* **5**:27-31, 1979
 - 23) Palevsky PM: Hyponatremia. *Semin Nephrol* **18**:20-30, 1998
 - 24) Blum D, Brasseur D, Kahn A, Brachet E: Safe oral rehydration of hypertonic dehydration. *J Pediatr Gastroenterol Nutr* **5**:232-235, 1986