

## Effects of Antioxidant Drugs in Rats with Acute Renal Injury

Eun Hui Bae, M.D.<sup>1</sup>, JongUn Lee, M.D.<sup>2</sup>, and Soo Wan Kim, M.D.<sup>1</sup>

Departments of <sup>1</sup>Internal Medicine and <sup>2</sup>Physiology, Chonnam National University Medical School, Gwangju, Korea

Acute renal failure is mainly caused by ischemia/reperfusion (I/R) injury or nephrotoxic drugs, in which reactive oxygen species (ROS) may play an important role. Therefore, antioxidants are expected to decrease the vulnerability of renal injury associated with oxidative challenges.  $\alpha$ -Lipoic acid ( $\alpha$ -LA), potent antioxidant, could act as ROS scavengers, iron chelators and enzyme modulators. In rats with acute renal injury, dysregulation of aquaporin (AQP) water channels and sodium transporters has been noted. I/R injury or cisplatin induced marked down-regulation of AQP1, AQP2 and AQP3 water channels, and type-3 Na-H exchanger, Na,K-ATPase, and Na-K-2Cl cotransporters, in association with impairment of urinary concentration and tubular sodium reabsorption. Treatment with  $\alpha$ -LA prevented the dysregulation of AQP channels and sodium transporters, along with improved urinary concentrating capability and renal sodium reabsorption.

**Key Words :** Aquaporins, Reperfusion injury, Sodium transporters, Thiocctic acid

### Introduction

Acute renal failure (ARF) is mainly caused by renal ischemia or nephrotoxic agents. It is characterized by severe reduction in glomerular filtration rate (GFR), urinary concentrating ability, and impaired tubular reabsorption of sodium. A diminished adenylylate cyclase activity and aquaporin-2 (AQP2) expression may contribute to the urinary concentration defect in post-ischemic kidneys<sup>1</sup>. Several studies reported defects in the collecting duct sodium handling in ischemia/reperfusion (I/R)-induced ARF rats<sup>2,3</sup>. Although the mechanisms underlying intrarenal vasoconstriction and hypoperfusion remain incompletely

defined, an imbalance of vasoactive agents such as endothelin and nitric oxide may be contributory<sup>4</sup>. On the other hand, the tubular cell damage may be caused by such factors as ATP depletion, intracellular calcium, oxygen free radicals and apoptosis<sup>5</sup>.

Under normal circumstances, the kidney generates reactive oxygen species (ROS), including superoxide anions, hydrogen peroxide, peroxynitrite and hydroxyl radical, which are efficiently eliminated by enzymatic-superoxide dismutase (SOD), catalase, glutathione peroxidase (GPX)-and non-enzymatic systems (glutathione, Vitamins C and E). Oxidative stress occurs when ROS production overrides metabolic capacity of the antioxidant defense system, often resulting in tissue damage (Fig. 1)<sup>6</sup>.

Numerous interventions have been put forward to counteract the effects of ROS, by reinforcing the antioxidant defense systems. Dietary supplementation with the antioxidant vitamin E slowed the rate of progression of renal deterioration<sup>7</sup>. Recently,  $\alpha$ -lipoic


Received March 7, 2007. Accepted May 10, 2007

Corresponding author: Soo Wan Kim, M.D.

Department of Internal Medicine, Chonnam National University Medical School, 8 Hak-dong, Dong-gu, Gwangju, 501-757, Korea

Tel: +82-62-2206272, Fax: +82-62-2258578

E-mail: kdksw@hanmail.net



**Fig. 1.** The balance of the oxidative stress and defence system.

Acid ( $\alpha$ -LA) has been known as a potent antioxidant (with a redox potential  $E_0^+$  of  $-290$  mV compared with vitamin E, which has a redox potential  $E_0^+$  of  $+370$  mV), endogenously existing in tissues and acting as a cofactor of key mitochondrial enzymes, controlling glucose oxidation, such as the pyruvate dehydrogenase and the  $\alpha$ -ketoglutarate dehydrogenase<sup>8</sup>. Its anti-oxidant effects are due to direct radical scavenging and metal chelation by both LA and its reduced form, dihydrolipoic acid (DHLA)<sup>9</sup>. It also has been demonstrated that LA reduces activation of nuclear factor (NF)- $\kappa$ b, a transcriptional factor in cultured endothelial cells and fractalkine-mediated inflammatory processes in endotoxemia<sup>10</sup>.

This review aims at providing recent insight into the role of  $\alpha$ -LA in the regulation of AQP water channels and sodium transporters in acute renal injury caused by I/R injury and cisplatin-treatment.

### Localization of AQP water channels and sodium (co)transporters

In the kidney, at least six isoforms of AQP proteins have been detected (AQP1, -2, -3, -4, -6, and -7). Among them, AQP1 is primarily responsible for the constitutive water permeability of proximal tubules and thin descending limbs of Henle<sup>11</sup>. Studies using AQP1-knockout mice revealed a dual role of AQP1 in proximal tubular reabsorption and medullary hypertonicity construction<sup>12</sup>. However, AQP2 to AQP4 are mainly expressed in the collecting duct. AQP2 is specifically located in the principal cells of the col-

lecting duct and is regulated in the short term and long term via the arginine vasopressin (AVP)/cyclic AMP pathway<sup>13</sup>. In addition, the constitutive localization of AQP3 and AQP4 in the basolateral membrane of principal cells confers to the epithelium high water permeability, in concert with AQP2 insertion into the apical membrane<sup>14</sup>.

The tubular reabsorption of solutes is basically linked to the activity of Na,K-ATPase that is heavily expressed in the basolateral membrane throughout the nephron segments. On the contrary, the sodium transporters in the apical membrane differ in different tubular segments. In the proximal tubule, type-3 Na-H exchanger (NHE3) is apically expressed. On the other hand, the apically expressed bumetanide-sensitive Na-K-2Cl cotransporters (NKCC2) and NHE3 are mainly responsible for the sodium reabsorption in the thick ascending loop of Henle (TAL). In the distal convoluted tubule, the thiazide-sensitive sodium chloride cotransporter (NCC) is involved in the apical movement of sodium<sup>14-16</sup>.

### Changes of AQP and sodium transporters in ischemia/reperfusion renal injury and cisplatin induced nephropathy

#### 1. Ischemia/reperfusion renal injury

The urinary concentration defect and increased water excretion is, at least in part, accounted for by a reduced abundance of AQP2 water channels in the collecting duct<sup>1</sup>. Furthermore, cAMP generation in response to AVP was attenuated in ARF, most

prominently in the outer medulla. Generation of cAMP in the outer medulla in response to forskolin was not affected, but sodium fluoride was significantly blunted in ARF. Therefore, the primary impairment may lie at the level of membrane G protein that acts as a transducer between AVP receptor and adenylate cyclase, decreasing the AVP-stimulated adenylate cyclase activity and hence subsequent generation of cAMP. We have also demonstrated that the abundance of several major renal sodium transporters (Na-K-ATPase, NHE-3, NKCC2, and NCC) was severely reduced following I/R injury, in association with an impairment of tubular reabsorption of filtered sodium. Decreased levels of sodium transporters along the nephron may play a role in the impairment of tubular sodium reabsorption.

Previous studies have shown that treatment with anti-inflammatory agents, significantly reduce the down-regulation of AQP-1,-2 and AQP-3 levels as well as major renal sodium transporters (NHE3, NaPi-2, BSC-1, TSC, and Na-K-ATPase) in rats with bilateral ischemia-induced ARF<sup>17, 18)</sup>. The anti-oxidant may play key role against the glomerular inflammatory processes, through a diminution of the activity of inflammatory enzymes and cytokine secretion, or by inhibiting the activity of NF- $\kappa$ B<sup>19)</sup>.

## 2. Cisplatin-induced nephropathy

Cisplatin is highly efficacious in treating a wide variety of neoplastic disease<sup>20)</sup>. However, its clinical use is often limited by its potential renal toxicity leading to ARF<sup>21-23)</sup>. The most silent and persistent features of cisplatin-induced nephropathy is polyuria and inability to concentrate urine<sup>24)</sup>. We have demonstrated that cisplatin-treatment caused polyuric renal failure in association with decrease in free water reabsorption<sup>25)</sup>. The expression of AQP1 and AQP2 was decreased in the cortex, outer medulla and inner medulla, whereas that of AQP3 was decreased in the outer medulla and the inner medulla. Immunohistochemistry also revealed that cisplatin decreased im-

munoreactivity for AQP1, AQP2, and AQP3. The AVP evoked generation of cAMP was attenuated by cisplatin, being most prominent in the outer medulla.

### Effects of $\alpha$ -LA on the dysregulation of AQP and sodium transporters

ARF was ameliorated by treatment with  $\alpha$ -LA at the time of ischemia<sup>26)</sup>. Accordingly, the creatinine clearance was increased. Furthermore,  $\alpha$ -LA treatment attenuated the polyuria that was encountered in the post-ischemic period, along with normally expressed abundance of AQP1-3. The reduced expression of NHE3 and Na,K-ATPase and the increase in  $FE_{Na}$  were completely prevented by treatment with  $\alpha$ -LA. Overall,  $\alpha$ -LA treatment 1) markedly inhibited the decline of GFR, 2) reduced the defects in urinary concentration and normalized the sodium handling, 3) prevented the down-regulation of AQP, being consistent with the improvement of urinary concentration ability and morphologic changes.

The underlying mechanism of  $\alpha$ -LA treatment to prevent I/R injury or cisplatin-induced nephropathy remains to be established. It was shown that nitric oxide inhibits AVP-stimulated osmotic water permeability in the isolated and perfused cortical collecting duct of the rat kidney<sup>27)</sup>. Yu et al<sup>28)</sup> demonstrated that nitric oxide production is stimulated by hypoxic injury in epithelial cells, and an inhibition of its synthesis protects against hypoxic injury in rat proximal tubule cells. Moreover, endothelin, and chemokines have been implicated in the urinary concentrating defect associated with ureteral obstruction, a condition which has been also associated with reduced expression of AQP1, AQP2, and AQP3<sup>29, 30)</sup>. These results suggest changes of renal hormones such as nitric oxide and endothelin may contribute to recovery renal function following  $\alpha$ -LA treatment in I/R- or cisplatin induced nephropathy. However, its underlying molecular mechanisms are probably multifactorial, and remain to be further elucidated.

## Conclusion

$\alpha$ -LA treatment has a protective effect against renal I/R injury and cisplatin-induced nephropathy in rat, such as normalization of urine output and  $FE_{Na}$  and regulation of AQP and sodium transporters.

## Acknowledgement

This study was supported by a grant (CUHRICM-Y-2007010) Chonnam national University Hospital Research Institute of Clinical Medicine.

## References

- 1) Kim SW, Jeon YS, Lee JU, Kang DG, Kook H, Ahn KY, Kim SZ, Cho KW, Kim NH, Han JS, Choi KC: Diminished adenylate cyclase activity and aquaporin 2 expression in acute renal failure rats. *Kidney Int* **57**:1643-1650, 2000
- 2) Johnston PA, Rennke H, Levinsky NG: Recovery of proximal tubular function from ischemic injury. *Am J Physiol* **246**:F159-F166, 1984
- 3) Tanner GA, Sloan KL, Sophasan S: Effects of renal artery occlusion on kidney function in the rat. *Kidney Int* **4**:377-389, 1973
- 4) Vetterlein F, Petho A, Schmidt G: Distribution of capillary blood flow in rat kidney during postischemic renal failure. *Am J Physiol* **251**:H510-H519, 1986
- 5) Vogt BA, Shanley TP, Croatt A, Alam J, Johnson KJ, Nath, KA: Glomerular inflammation induces resistance to tubular injury in the rat. A novel form of acquired, heme oxygenase-dependent resistance to renal injury. *J Clin Invest* **98**:2139-2145, 1996
- 6) Vaya J, Aviram M: Nutritional antioxidant: mechanisms of action, analyses of activities and medical applications. *Curr Med Chem-Imm Endoc & Metab Agents* **1**:99-117, 2001
- 7) Fryer MJ: Vitamin E may slow kidney failure owing to oxidative stress. *Redox Rep* **3**:259-261, 1997
- 8) Packer L, Roy S, Sen CK:  $\alpha$ -lipoic acid: a metabolic antioxidant and potential redox modulator of transpiration. *Adv Pharmacol* **38**:79-101, 1997
- 9) Packer L, Witt EH, Tritschler HJ:  $\alpha$ -Lipoic acid as a biological antioxidant. *Free Radic Biol Med* **19**:227-250, 1995
- 10) Sung MJ, Kim W, Ahn SY, Cho CH, Koh GY, Moon SO, Kim DH, Lee S, Kang KP, Jang KY, Park SK: Protective effect of  $\alpha$ -lipoic acid in lipopolysaccharide-induced endothelial fractalkine expression. *Circ Res* **97**:880-890, 2005
- 11) Sabolic I, Valenti G, Verbavatz JM, Van Hoek AN, Verkman AS, Ausiello DA, Brown D: Localization of the CHIP28 water channel in rat kidney. *Am J Physiol* **263**:C1225-C1233, 1992
- 12) Ma T, Yang B, Gillespie A, Carlson EJ, Epstein CJ, Verkman AS: Severely impaired urinary concentrating ability in transgenic mice lacking aquaporin-1 water channels. *J Biol Chem* **273**:4296-4299, 1998
- 13) Yamamoto T, Sasaki S, Fushimi K, Ishibashi K, Yaoita E, Kawasaki K, Marumo F, Kihara I: Vasopressin increases AQP-CD water channel in apical membrane of collecting duct cells in Brattleboro rats. *Am J Physiol* **268**:C1546-C1551, 1995
- 14) Kashgarian M, Biemesderfer D, Caplan M, Forbush B 3rd: Monoclonal antibody to Na,K-ATPase: immunocytochemical localization along nephron segments. *Kidney Int* **28**:899-913, 1985
- 15) Kim SW, Wang W, Kwon TH, Knepper MA, Frokiaer J, Nielsen S: Increased expression of ENaC subunits and increased apical targeting of AQP2 in the kidneys of spontaneously hypertensive rats. *Am J Physiol Renal Physiol* **289**:F957-968, 2005
- 16) Kim SW, Wang W, Nielsen J, Praetorius J, Kwon TH, Knepper MA, Frokiaer J, Nielsen S: Increased expression and apical targeting of renal ENaC subunits in puromycin aminonucleoside-induced nephrotic syndrome in rats. *Am J Physiol Renal Physiol* **286**:F922-F935, 2004
- 17) Kwon TH, Frokiaer J, Han JS, Knepper MA, Nielsen S: Decreased abundance of major Na(+) transporters in kidneys of rats with ischemia-induced acute renal failure. *Am J Physiol Renal Physiol* **278**:F925-F939, 2000
- 18) Kwon TH, Frokiaer J, Fernandez-Llama P, Knepper MA, Nielsen S: Reduced abundance of aquaporins in rats with bilateral ischemia-induced acute renal failure: prevention by  $\alpha$ -MSH. *Am J Physiol* **277**:F413-F427, 1999
- 19) Massy ZA, Guijarro C, O'Donnell MP, Kim Y, Kashtan CE, Egido J, Kasiske BL, Keane WF: The central role of nuclear factor- $\kappa$ B in mesangial cell activation. *Kidney Int* **56**(Suppl.71):S76-S79, 1999
- 20) Lebwohl D, Canetta R: Clinical development of platinum complexes in cancer therapy: an historical perspective and an update. *Eur J Cancer* **34**:1522-1534, 1998
- 21) Goldstein RS, Mayor GH: Minireview. The nephrotoxicity of cisplatin. *Life Sci* **32**:685-690, 1983
- 22) Choie DD, Longnecker DS, del Campo AA: Acute and chronic cisplatin nephropathy in rats. *Lab Invest* **44**:397-402, 1981

- 23) Dobyas DC, Levi J, Jacobs C, Kosek J, Weiner MW : Mechanism of cis-platinum nephrotoxicity: II. Morphologic observations. *J Pharmacol Exp Ther* **213**:551-556, 1980
- 24) Ecelbarger CA, Sands JM, Doran JJ, Cacini W, Kishore BK : Expression of salt and urea transporters in rat kidney during cisplatin-induced polyuria. *Kidney Int* **60**:2274-2282, 2001
- 25) Kim SW, Lee JU, Nah MY, Kang DG, Ahn KY, Lee HS, Choi KC : Cisplatin decreases the abundance of aquaporin water channels in rat kidney. *J Am Soc Nephrol* **12**:875-882, 2001
- 26) Takaoka M, Ohkita M, Kobayashi Y, Yuba M, Matsumura Y : Protective effect of alpha-lipoic acid against ischaemic acute renal failure in rats. *Clin Exp Pharmacol Physiol* **29**:189-194, 2002
- 27) Garcia NH, Pomposiello SI, Garvin JL : Nitric oxide inhibits ADH-stimulated osmotic water permeability in cortical collecting ducts. *Am J Physiol* **270**:F206-F210, 1996
- 28) Yu L, Gengaro PE, Niederberger M, Burke TJ, Schrier RW : Nitric oxide: a mediator in rat tubular hypoxia/reoxygenation injury. *Proc Natl Acad Sci U S A* **91**:1691-1695, 1994
- 29) Frokiaer J, Marples D, Knepper MA, Nielsen S : Bilateral ureteral obstruction downregulates expression of vasopressin-sensitive AQP-2 water channel in rat kidney. *Am J Physiol* **270**:F657-F668, 1996
- 30) Frokiaer J, Christensen BM, Marples D, Djurhuus JC, Jensen UB, Knepper MA, Nielsen S : Downregulation of aquaporin-2 parallels changes in renal water excretion in unilateral ureteral obstruction. *Am J Physiol* **273**:F213-F223, 1997