

Renal Handling of Ammonium and Acid Base Regulation

Hye-Young Kim, M.D.

Department of Internal Medicine, Chungbuk National University College of Medicine, Cheongju, Korea

Renal ammonium metabolism is the primary component of net acid excretion and thereby is critical for acid-base homeostasis. Briefly, ammonium is produced from glutamine in the proximal tubule in a series of biochemical reactions that result in equimolar bicarbonate. Ammonium is predominantly secreted into the luminal fluid via the apical Na^+/H^+ exchanger, NHE3. The thick ascending limb of the loop of Henle reabsorbs luminal ammonium, predominantly by transport of NH_4^+ by the apical $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporter, BSC1/NKCC2. This process results in renal interstitial ammonium accumulation. Finally, the collecting duct secretes ammonium from the renal interstitium into the luminal fluid. Although in past ammonium was believed to move across epithelia entirely by passive diffusion, an increasing number of studies demonstrated that specific proteins contribute to renal ammonium transport. Recent studies have yielded important new insights into the mechanisms of renal ammonium transport. In this review, we will discuss renal handling of ammonium, with particular emphasis on the transporters involved in this process.

Key Words : ammonia; kidney; kidney tubules, collecting; acidosis

Introduction

The acid-base regulation is chiefly dependent on the control of net acid excretion by the kidney and CO_2 excretion by the lungs. Renal acid-base homeostasis consists of two major processes, the reabsorption of filtered bicarbonate and the excretion of the hydrogen ion. The kidney excretes hydrogen ion through the processes of titratable acid excretion and urinary ammonium excretion^{1, 2)}. Quantitatively, urinary ammonium excretion is the primary mechanism of net acid excretion both under basal conditions and in response to acid loads¹⁻³⁾. The understanding renal ammonium production and transport is of fundamental importance for understanding acid-base homeostasis.

The term "total ammonia" is used to denote the sum of NH_3 and NH_4^+ . In this review, because the vast major-

ity of the total ammonia is in the form of NH_4^+ at the physiologic pH, we generally refer to total ammonia transport as "ammonium transport" and to total ammonia excretion as "ammonium excretion"⁴⁾.

Renal ammonium production

The proximal tubule is the chief site of renal ammonium production. Glutamine catabolism in the proximal tubule generates NH_4^+ and also bicarbonate after complete catabolism of α -ketoglutarate to CO_2 and H_2O . Studies in microdissected tubules have demonstrated that proximal tubules from the acidotic rats produced substantially more ammonium than did tubules from controls⁵⁾. These results illustrate that the production of ammonium is regulated according to the acid-base state. In addition, hormonal factors contribute to the stimulation by metabolic acidosis of ammoniogenesis in the proximal tubule. The circulating levels of glucocorticoids increase during metabolic acidosis⁶⁾.

Renal ammonium transport

Renal ammonium metabolism and transport involves

Received May 18, 2009. Accepted May 27, 2009.

Corresponding author : Hye-Young Kim, M.D.

Department of Internal Medicine, Chungbuk National University College of Medicine, 62 Gaeshin-dong, Heunduk-gu, Cheongju, 361-711, Korea
Tel : +82-43-269-6017, Fax : +82-43-273-3252

E-mail : hyekim@chungbuk.ac.kr

integrated responses of multiple portions of the kidney, including specific transport mechanisms in the proximal tubule, thick ascending limb of the loop of Henle, and the collecting duct.

1. Ammonium transport in the proximal tubule

The proximal tubule secretes NH_4^+ into the luminal fluid primarily through the action of the apical sodium/hydrogen ion exchanger, NHE3^{7, 8)}, and to a lesser degree through an apical barium sensitive potassium ion channel⁸⁻¹⁰⁾. The activity and abundance of the apical Na^+/H^+ exchanger NHE3 is increased in the proximal brush border membrane during metabolic acidosis, which could be expected to contribute to the enhanced NH_4^+ secretion by the proximal tubule during this condition¹¹⁾. It was recently shown that stimulation of ammonium secretion by the proximal tubule during metabolic acidosis largely depends on $\text{Na}^+/\text{H}^+(\text{NH}_4^+)$ exchange activity and on angiotensin II¹²⁾.

2. Ammonium transport in thick ascending limb of Henle's loop

After ammonium is produced and secreted by the proximal tubule, it is then delivered into the renal medulla via the loop of Henle. These potential losses of luminal NH_3 are minimized because more than 75 percent of the tubular fluid NH_4^+ is recycled within the medulla, thereby maintaining a high interstitial NH_3 concentration^{3, 13, 14)}. The primary step in this process is reabsorption in the thick ascending limb by substitution of NH_4^+ for K^+ both on the $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ carrier and, to a much lesser degree, through the K^+ channels in the luminal membrane^{3, 15)}. The luminal NH_3 permeability in the thick ascending limb was lower compared to other nephron segments¹⁶⁾. The low NH_3 permeability limits the backflux of NH_3 into the tubule lumen and thereby contributes to the overall efficiency of the NH_4^+ absorptive process. Partial dissociation into NH_3 and H^+ then occurs in the less acidic tubular cell. As a result, the NH_3 formed within the cell will diffuse out across the basolateral membrane into the medullary interstitium. The countercurrent multiplication of ammonium generates the maintenance of a high medullary

interstitial NH_3 concentration which promotes secretion into the medullary collecting tubule.

Ammonium reabsorption in the thick ascending limb of Henle's loop is reduced by hyperkalemia and is enhanced by chronic metabolic acidosis due to increased NH_4^+ production in and delivery out of the proximal tubule^{14, 17)}. In the thick ascending limb of the loop of Henle, metabolic acidosis increases ammonium reabsorption through mechanisms that appear to be involved in increasing NKCC2 expression^{17, 18)}. In vitro incubation of rat medullary thick ascending limb fragments in suspension in an acid medium strongly enhanced the BSC1/NKCC2 mRNA and protein abundance and cotransport activity¹⁸⁾. In addition, administration of the glucocorticoid dexamethasone to adrenalectomized rats stimulated BSC1/NKCC2 expression at the mRNA and protein levels¹⁹⁾. The acidity of the surrounding environment and glucocorticoids may account for the stimulating effect of chronic metabolic acidosis on BSC1/NKCC2 expression.

3. Ammonium secretion in the collecting duct

The majority of urinary ammonium is secreted into luminal fluid in the region of the nephron distal to the micropuncturable late distal tubule²⁰⁾. This is a heterogeneous region, and includes portions of the distal convoluted tubule (DCT), connecting segment (CNT), initial collecting tubule, cortical collecting duct (CCD), outer medullary collecting duct (OMCD) and inner medullary collecting duct (IMCD). Accordingly, understanding the mechanisms of transepithelial ammonium transport across the cells that comprise these portions of the kidney is important.

The mechanism of ammonium secretion in OMCD and IMCD is not completely clear at present. Since the first description of the concept, the process of transepithelial transport of ammonium in the collecting duct is thought to occur primarily through passive non-ionic NH_3 diffusion^{21, 22)}. The fluid entering the collecting tubules has a relatively low NH_3 concentration, because of its removal in the loop of Henle. The net effect is that there is a relatively large gradient favoring the free diffusion of inter-




Fig. 1. Schematic representation of the ammonium transport mechanisms in the kidney. Ammonium is predominantly secreted into the luminal fluid via the apical Na^+/H^+ exchanger, NHE3. The thick ascending limb of the loop of Henle reabsorbs luminal ammonium, predominantly by transport of NH_4^+ by the apical $\text{Na}^+/\text{K}^+/\text{2Cl}^-$ cotransporter, BSC1/NKCC2. The renal countercurrent mechanism results in renal interstitial ammonium accumulation. Finally, the collecting duct secretes ammonium from the renal interstitium into the luminal fluid. NH_3 is transported across the basolateral membrane, predominantly by Rhcg, but also by lipid diffusion (dashed line) and possibly by Rhbg. Intracellular NH_3 is secreted across the apical membrane by apical Rhcg. In inner medullary collecting duct, basolateral Na^+/K^+ -ATPase transports NH_4^+ . There is also likely to be a component of diffusive apical NH_3 transport (dashed line). H^+ -ATPase secretes H^+ , which combines with luminal NH_3 to form NH_4^+ . Modified from Kim et al.⁴².

stitial NH_3 into the tubular lumen, where it forms NH_4^+ ^{1, 13}. The cell membranes in the collecting tubules are highly permeable to NH_3 but have only a negligible permeability to NH_4^+ ²³. As a result, interstitial NH_3 can passively diffuse into the tubular lumen where it is then trapped as NH_4^+ . The net effect is that NH_3 is secreted into the lumen throughout the collecting tubules³.

Although in the past ammonium was believed to move across epithelia entirely by passive diffusion, an increasing number of studies demonstrate that specific proteins contribute to renal ammonium transport^{3, 24, 25}. The basolateral $\text{Na}^+/\text{K}^+(\text{NH}_4^+)\text{-ATPase}$ pump has been demonstrated to mediate NH_4^+ secretion in the rat IMCD isolated and perfused *in vitro*²⁶. The secretory $\text{Na}^+/\text{K}^+(\text{NH}_4^+)/\text{2Cl}^-$ cotransporter BSC2/NKCC1 was recently reported to mediate K^+ - and NH_4^+ - dependent chloride secretion, and thus to be involved in transepithelial solute transport²⁷. BSC2/NKCC1 was shown to be up-regulated by chronic metabolic acidosis in collecting ducts of the rat²⁸.

The most recent addition to our understanding of the molecular mechanisms of ammonium metabolism is the identification of the ammonia transporter family of proteins^{25, 29}. Rh B glycoprotein (Rhbg) and Rh C glycoprotein (Rhcg) are expressed in the renal DCT, CNT and collecting duct, the sites where approximately 80% of urinary ammonium is secreted³⁰⁻³⁶. In conditions of increased single-nephron ammonium metabolism, such as metabolic acidosis and reduced renal mass, Rhcg expression increases, suggesting that Rhcg mediates and has an important role in renal ammonium transport^{32, 35, 36}. However, the nature of the transported substrate (NH_4^+ or NH_3) by Rh glycoprotein has been controversial³⁷⁻³⁹. It was recently shown that both global and collecting duct-specific Rhcg deletion altered renal urinary ammonium excretion, indicating that collecting duct ammonium secretion is, at least in part, mediated by Rhcg and not solely by lipid diffusion^{40, 41}. In addition, *in vitro* microperfused collecting ducts of Rhcg^{-/-} acid-loaded mice show reduced ap-

ical permeability to NH_3 and impaired transepithelial NH_3 transport, indicating a role of Rhcg as an ammonium transport protein mediating the net flux of NH_3 in the collecting duct.

Both the rat and the human kidney express both apical and basolateral Rhcg immunolabel^{30, 32, 35, 36}. Basolateral Rhcg expression increases in parallel with urinary ammonium excretion in both metabolic acidosis and reduced renal mass, suggesting that changes in basolateral Rhcg expression may contribute to renal ammonium metabolism^{35, 36}. We recently demonstrate that the mouse kidney expresses both apical and basolateral Rhcg, which is similar to findings in both the rat and human kidney, although there are strain-dependent differences in the level of basolateral Rhcg expression in the mouse kidney⁴². These observations suggest that basolateral Rhcg may contribute to facilitated transcellular ammonium movement. Furthermore, basolateral Rhcg is likely to be the same protein as apical Rhcg, but is trafficked to the basolateral plasma membrane. Fig. 1 summarizes the major transporters involved in renal ammonium transport.

In summary, renal ammonium handling involves intrarenal ammonium production and transepithelial transport in multiple tubular segments that results in highly regulated renal ammonium metabolism. An increasing number of studies demonstrate that specific proteins contribute to renal ammonium transport. Understanding the physiologic regulation of ammonium transporter and the contribution of other protein to renal ammonium transport, are likely to be important fields for future studies.

Acknowledgements

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, Basic Research Promotion Fund) (KRF-2008-E00035).

References

- Hamm LL, Simon EE: Roles and mechanisms of urinary buffer excretion. *Am J Physiol* 253:F595-605, 1987
- Halperin ML, Jungas RL: Metabolic production and renal disposal of hydrogen ions. *Kidney Int* 24:709-713, 1983
- DuBose TD Jr, Good DW, Hamm LL, Wall SM: Ammonium transport in the kidney: new physiological concepts and their clinical implications. *J Am Soc Nephrol* 1:1193-1203, 1991
- Knepper MA: NH_4^+ transport in the kidney. *Kidney Int Suppl* 40:S95-102, 1991
- Good DW, Burg MB: Ammonia production by individual segments of the rat nephron. *J Clin Invest* 73:602-610, 1984
- May RC, Kelly RA, Mitch WE: Metabolic acidosis stimulates protein degradation in rat muscle by a glucocorticoid-dependent mechanism. *J Clin Invest* 77:614-621, 1986
- Nagami GT: Ammonia production and secretion by the proximal tubule. *Am J Kidney Dis* 14:258-261, 1989
- Simon EE, Merli C, Herndon J, Cragoe EJ Jr, Hamm LL: Effects of barium and 5-(N-ethyl-N-isopropyl)-amiloride on proximal tubule ammonia transport. *Am J Physiol* 262:F36-39, 1992
- Hamm LL, Simon EE: Ammonia transport in the proximal tubule. *Miner Electrolyte Metab* 16:283-290, 1990
- DuBose TD Jr, Good DW: Effects of chronic hyperkalemia on renal production and proximal tubule transport of ammonium in rats. *Am J Physiol* 260:F680-687, 1991
- Nagami GT: Luminal secretion of ammonia in the mouse proximal tubule perfused in vitro. *J Clin Invest* 81:159-164, 1988
- Nagami GT: Ammonia production and secretion by S3 proximal tubule segments from acidotic mice: role of ANG II. *Am J Physiol Renal Physiol* 287:F707-712, 2004
- Buerkert J, Martin D, Trigg D: Segmental analysis of the renal tubule in buffer production and net acid formation. *Am J Physiol* 244:F442-454, 1983
- Packer RK, Desai SS, Hornbuckle K, Knepper MA: Role of countercurrent multiplication in renal ammonium handling: regulation of medullary ammonium accumulation. *J Am Soc Nephrol* 2:77-83, 1991
- Garvin JL, Burg MB, Knepper MA: Active NH_4^+ absorption by the thick ascending limb. *Am J Physiol* 255:F57-65, 1988
- Kikeri D, Sun A, Zeidel ML, Hebert SC: Cell membranes impermeable to NH_3 . *Nature* 339:478-480, 1989
- Good DW: Adaptation of HCO_3^- and NH_4^+ transport in rat MTAL: effects of chronic metabolic acidosis and Na^+ intake. *Am J Physiol* 258:F1345-1353, 1990
- Attmane-Elakeb A, Mount DB, Sibella V, Vernimmen C, Hebert SC, Bichara M: Stimulation by in vivo and in vitro metabolic acidosis of expression of rBSC-1, the $\text{Na}^+\text{-K}^+$ (NH_4^+)- 2Cl^- cotransporter of the rat medullary thick ascending limb. *J Biol Chem* 273:33681-33691, 1998
- Wall SM: NH_4^+ augments net acid secretion by a ouabain-sensitive mechanism in isolated perfused inner medullary collecting ducts. *Am J Physiol* 270:F432-439, 1996
- Sajo IM, Goldstein MB, Sonnenberg H, Stinebaugh BJ, Wilson DR, Halperin ML: Sites of ammonia addition to tubular fluid in rats with chronic metabolic acidosis. *Kidney*

- Int 20:353-358, 1981
- 21) Pitts RF: Renal excretion of acid. *Fed Proc* 7:418-426, 1948
 - 22) Pitts RF: The role of ammonia production and excretion in regulation of acid-base balance. *N Engl J Med* 284:32-38, 1971
 - 23) Flessner MF, Wall SM, Knepper MA: Permeabilities of rat collecting duct segments to NH_3 and NH_4^+ . *Am J Physiol* 260:F264-272, 1991
 - 24) Wall SM: Ammonium transport and the role of the Na,K-ATPase. *Miner Electrolyte Metab* 22:311-317, 1996
 - 25) Weiner ID, Hamm LL: Molecular mechanisms of renal ammonia transport. *Annu Rev Physiol* 69:317-340, 2007
 - 26) Wall SM, Koger LM: NH_4^+ transport mediated by Na(+)-K(+)-ATPase in rat inner medullary collecting duct. *Am J Physiol* 267:F660-670, 1994
 - 27) Wall SM, Fischer MP, Mehta P, Hassell KA, Park SJ: Contribution of the $\text{Na}^+\text{-K}^+\text{-2Cl}^-$ cotransporter NKCC1 to Cl^- secretion in rat OMCD. *Am J Physiol Renal Physiol* 280:F913-921, 2001
 - 28) Ikebe M, Nonoguchi H, Nakayama Y, Tashima Y, Tomita K: Upregulation of the secretory-type Na(+)/K(+)/2Cl(-)-cotransporter in the kidney by metabolic acidosis and dehydration in rats. *J Am Soc Nephrol* 12:423-430, 2001
 - 29) Heitman J, Agre P: A new face of the Rhesus antigen. *Nat Genet* 26:258-259, 2000
 - 30) Han KH, Croker BP, Clapp WL, Werner D, Sahni M, Kim J, et al.: Expression of the ammonia transporter, Rh C glycoprotein, in normal and neoplastic human kidney. *J Am Soc Nephrol* 17:2670-2679, 2006
 - 31) Quentin F, Eladari D, Cheval L, Lopez C, Goossens D, Colin Y, et al.: RhBG and RhCG, the putative ammonia transporters, are expressed in the same cells in the distal nephron. *J Am Soc Nephrol* 14:545-554, 2003
 - 32) Seshadri RM, Klein JD, Kozlowski S, Sands JM, Kim YH, Han KH, et al.: Renal expression of the ammonia transporters, Rhbg and Rhcg, in response to chronic metabolic acidosis. *Am J Physiol Renal Physiol* 290:F397-408, 2006
 - 33) Verlander JW, Miller RT, Frank AE, Royaux IE, Kim YH, Weiner ID: Localization of the ammonium transporter proteins RhBG and RhCG in mouse kidney. *Am J Physiol Renal Physiol* 284:F323-337, 2003
 - 34) Weiner ID, Verlander JW: Renal and hepatic expression of the ammonium transporter proteins, Rh B Glycoprotein and Rh C Glycoprotein. *Acta Physiol Scand* 179:331-338, 2003
 - 35) Kim HY, Baylis C, Verlander JW, Han KH, Reungjui S, Handlogten ME, et al.: Effect of reduced renal mass on renal ammonia transporter family, Rh C glycoprotein and Rh B glycoprotein, expression. *Am J Physiol Renal Physiol* 293:F1238-1247, 2007
 - 36) Seshadri RM, Klein JD, Smith T, Sands JM, Handlogten ME, Verlander JW, et al.: Changes in subcellular distribution of the ammonia transporter, Rhcg, in response to chronic metabolic acidosis. *Am J Physiol Renal Physiol* 290:F1443-1452, 2006
 - 37) Nakhoul NL, Dejong H, Abdunour-Nakhoul SM, Boulpaep EL, Hering-Smith K, Hamm LL: Characteristics of renal Rhbg as an NH_4^+ transporter. *Am J Physiol Renal Physiol* 288:F170-181, 2005
 - 38) Nakhoul NL, Schmidt E, Abdunour-Nakhoul SM, Hamm LL: Electrogenic ammonium transport by renal Rhbg. *Transfus Clin Biol* 13:147-153, 2006
 - 39) Mak DO, Dang B, Weiner ID, Foskett JK, Westhoff CM: Characterization of ammonia transport by the kidney Rh glycoproteins RhBG and RhCG. *Am J Physiol Renal Physiol* 290:F297-305, 2006
 - 40) Biver S, Belge H, Bourgeois S, Van Vooren P, Nowik M, Scohy S, et al.: A role for Rhesus factor Rhcg in renal ammonium excretion and male fertility. *Nature* 456:339-343, 2008
 - 41) Lee HW, Verlander JW, Bishop JM, Igarashi P, Handlogten ME, Weiner ID: Collecting duct-specific Rh C glycoprotein deletion alters basal and acidosis-stimulated renal ammonia excretion. *Am J Physiol Renal Physiol* 296:F1364-1375, 2009
 - 42) Kim HY, Verlander JW, Bishop JM, Cain BD, Han KH, Igarashi P, et al.: Basolateral expression of the ammonia transporter family member Rh C glycoprotein in the mouse kidney. *Am J Physiol Renal Physiol* 296:F543-555, 2009