

The Role of Proximal Nephron in Cyclophosphamide-Induced Water Retention: Preliminary Data

Sua Kim, M.S.¹
Chor Ho Jo, M.S.¹
Joon-Sung Park, M.D.¹
Ho Jae Han, Ph.D.²
Gheun-Ho Kim, M.D.¹

¹Department of Internal Medicine, Hanyang University College of Medicine, Seoul, Korea

²Department of Veterinary, Chonnam National University College of Veterinary Medicine, Gwangju, Korea

Received: June 13, 2011

Accepted: June 21, 2011

Corresponding author: Gheun-Ho Kim, M.D., Ph.D.
Department of Internal Medicine, Hanyang University College of Medicine, 17 Haengdang-dong, Seongdong-gu, Seoul, 133-792, Korea
Tel: +82-2-2290-8318, Fax: +82-2-2298-9183
E-mail: kimgh@hanyang.ac.kr

Cyclophosphamide is clinically useful in treating malignancy and rheumatologic disease, but has limitations in that it induces hyponatremia. The mechanisms by which cyclophosphamide induces water retention in the kidney have yet to be identified. This study was undertaken to test the hypothesis that cyclophosphamide may produce water retention via the proximal nephron, where aquaporin-1 (AQP1) and aquaporin-7 (AQP7) water channels participate in water absorption. To test this hypothesis, we gave a single dose of intraperitoneal cyclophosphamide to male Sprague-Dawley rats and treated rabbit proximal tubule cells (PTCs) with 4-hydroperoxycyclophosphamide (4-HC), an active metabolite of cyclophosphamide. In the short-term 3-day rat study, AQP1 protein expression was significantly increased in the whole kidney homogenates by cyclophosphamide administration at 48 ($614 \pm 194\%$, $P < 0.005$), and 96 ($460 \pm 46\%$, $P < 0.05$) mg/kg BW compared with vehicle-treated controls. Plasma sodium concentration was significantly decreased (143 ± 1 vs. 146 ± 1 mEq/L, $P < 0.05$) by cyclophosphamide 100 mg/kg BW in the long-term 6-day rat study. When primary cultured rabbit PTCs were treated with 4-HC for 24 hours, the protein expressions of AQP1 and AQP7 were increased in a dose-dependent manner. Quantitative polymerase chain reaction revealed no significant changes in the mRNA levels of AQP1 and AQP7 from cyclophosphamide-treated rat renal cortices. From these preliminary data, we conclude that the proximal nephron may be involved in cyclophosphamide-induced water retention via AQP1 and AQP7 water channels. Further studies are required to demonstrate intracellular mechanisms that affect the expression of AQP proteins.

Key Words: cyclophosphamide; aquaporin 1; hyponatremia; water; proximal tubule

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Some pharmacologic agents have been shown to have water-retaining properties. Whereas the antidiuretic actions are used to advantage in the treatment of patients with diabetes insipidus, there have been numerous reports

of serious water retention and dilutional hyponatremia as a consequence of the water-retaining drugs¹. Thus, hyponatremia is one of the common electrolyte abnormalities induced by drugs.

Cyclophosphamide is an alkylating agent that is useful for the treatment of malignancy and rheumatologic

disease. It can produce an antidiuretic effect when injected intravenously into water-loaded patients^{2,3}. We recently reported that cyclophosphamide-induced hyponatremia is not rare even when low-dose pulse cyclophosphamide is used⁴. However, the mechanisms by which cyclophosphamide induces water retention in the kidney have not been defined.

The main water channels participating in water conservation along the renal tubule are aquaporin-1 (AQP1) and aquaporin-2 (AQP2). AQP1 is abundant in the apical and basolateral membranes of renal proximal tubules and descending thin limbs. AQP2 is localized in the apical membrane and in intracellular vesicles which are targeted to the apical plasma membranes when stimulated by vasopressin⁵. The traditional view was that some drugs including cyclophosphamide may potentiate the action of vasopressin on the kidney, but this explanation has never been verified in cases with cyclophosphamide-induced water retention.

In this study, we hypothesized that in producing water retention in the proximal tubule and Henle's thin descending limb, the AQP1 water channel may have a role in cyclophosphamide-induced hyponatremia. To test this hypothesis, the expression of AQP1 was investigated both *in vivo* and *in vitro* after the administration of cyclophosphamide and its active metabolite 4-hydroperoxycyclophosphamide (4-HC), respectively.

Materials and Methods

1. Animal experiments of cyclophosphamide administration

Specific pathogen-free male Sprague-Dawley rats (Orient Bio Inc., Seongnam, Korea) weighing 240–260 g, were used for two animal experiments of single cyclophosphamide administration. For the short-term 3-day study (Animal Experiment I), different doses (12, 24, 48, and 96 mg/kg BW) of intraperitoneal cyclophosphamide were administered to rats in Animal Experiment I. Control rats

received an injection of the vehicle (normal saline) solution only. All rats were provided with a gelled agar (1%) diet, so that adequate water loading (52 mL/200 g BW per day) and a fixed amount of daily regular rat chow (18 g/200 g BW per day) were given to each rat. In addition, they were freely accessible to drinking water.

For the long-term 6-day study (Animal Experiment II), a large dose (100 mg/kg BW) of intraperitoneal cyclophosphamide was given to rats in Animal Experiment II. For water loading, drinking water containing 600 mM sucrose was supplied. Rats spontaneously drink large amounts of this fluid, causing a water diuresis⁶. Serum and urine samples were collected at the time each rat was killed for determination of electrolytes and osmolality. The experimental protocols were approved by the institutional Animal Care and Use Committee of Hanyang University.

2. Proximal tubular cell culture and treatment with 4-HC

Primary cultured renal proximal tubular cells (PTCs) of New Zealand White male rabbits (1.5–2.0 kg) from Dae Han Experimental Animal (Chungju, Korea) were prepared as previously described⁷. The PTCs were grown in DMEM/F-12 medium supplemented with 15 mM HEPES and 20 mM sodium bicarbonate (pH 7.4). Three other growth supplements (5 µg/mL insulin, 5 µg/mL transferrin, and 5×10^{-8} M hydrocortisone) were added immediately before the medium was used. The kidneys from a rabbit were perfused through the renal artery, first with phosphate-buffered saline (PBS) and then with medium containing 0.5% iron oxide. Renal cortical slices were prepared and homogenized. The homogenate was poured sequentially through 253- and 83-µm mesh filters. Tubules and glomeruli caught by the 83-µm filter were transferred into sterile medium. The glomeruli (which contained the iron oxide) were removed using a magnetic stir bar. The remaining proximal tubules were incubated briefly in a medium containing 0.125 mg/mL collagenase and 0.025% soybean trypsin inhibitor. The tubules were then washed by centrifugation, resuspended in a medium containing the three supplements, and transferred into

tissue culture dishes. The medium was changed 1 day after plating and every 2 days thereafter. The primary cultured rabbit kidney PTCs were maintained at 37°C in a 5% CO₂ humidified environment and a serum-free basal medium amended with the 3 growth supplements.

4-HC was obtained from Niomec (Bielefeld, Germany). According to the previous study applying on the human hair follicle⁸⁾, 4 different 4-HC concentrations (1, 3, 10, and 30 μ mol/L) were prepared on ice in Williams' E medium immediately before use because of the short half-life of the compound⁹⁾. These concentrations were previously selected to imitate relevant serum levels of 4-HC in patients receiving moderate- to high-dose cyclophosphamide therapy¹⁰⁻¹²⁾. Cells were harvested in 1 to 3 days after the 4-HC treatment for immunoblot analysis.

3. Preparation of cytosolic and membrane fraction from PTCs

Once confluent cultures were treated according to the indicated conditions, the media was removed and PTCs were washed twice with ice-cold PBS, scraped, harvested by microcentrifugation, and resuspended in Buffer A (137 mM NaCl, 8.1 mM Na₂HPO₄, 2.7 mM KCl, 1.5 mM KH₂PO₄, 2.5 mM EDTA, 1 mM dithiothreitol, 0.1 mM phenylmethanesulfonyl fluoride, 10 μ g/mL leupeptin [pH 7.5]). Resuspended cells were then lysed mechanically on ice by trituration with a 21.1-gauge needle. To prepare the cytosolic and particulate fractions, the lysates were centrifuged at 1,000 μ g for 10 minutes at 4°C, after which the supernatants were centrifuged at 100,000 μ g for 1 h at 4°C. The particulate fraction, which contained the membranes, was washed twice and resuspended in Buffer A containing 1% (v/v) Triton X-100. Protein levels in each fraction were quantified using the Bradford procedure¹³⁾.

4. Immunoblot analysis

When each animal experiment was finished, whole kidneys or manually dissected slices of renal cortex were homogenized in a buffer containing 250 mM sucrose, 10 mM triethanolamine, 1 μ g/mL leupeptin, and 0.1 mg/

mL phenylmethanesulfonyl fluoride titrated to pH 7.6. Coomassie-stained "loading gels" were done to assess the quality of the protein by sharpness of the bands and to adjust protein concentrations before immunoblotting¹⁴⁾. For immunoblotting, the proteins were transferred electrophoretically from unstained gels to nitrocellulose membranes (Bio-Rad, Hercules, CA, USA). After being blocked with 5% skim milk in PBS-T (80 mM Na₂HPO₄, 20 mM NaH₂PO₄, 100 mM NaCl, 0.1 percent Tween-20, pH 7.5) for 30 minutes, membranes were probed overnight at 4°C with the respective primary antibodies. The secondary antibody was goat anti-rabbit IgG conjugated to horseradish peroxidase (Pierce, Rockford, IL, USA). Sites of antibody-antigen reaction were viewed using enhanced chemiluminescence substrate (ECLTM RPN 2106; Amersham Pharmacia Biotech, Buckinghamshire, UK) before exposure to X-ray film (Hyperfilm; Amersham Pharmacia Biotech, Buckinghamshire, UK). Relative quantification of the band densities from immunoblots was carried out by densitometry using a laser scanner and ImageQuaNT software (Molecular Analyst version 1.5, Bio-Rad, Hercules, CA, USA).

5. Quantitative polymerase chain reaction (QPCR)

Total RNA was isolated from the rat whole kidneys with the TRIzol[®] Reagent (Life Technologies, Carlsbad, CA, USA). RNA was quantified by spectrophotometry, and cDNA synthesis was performed on 3 μ g of RNA with SuperScript[®] III Reverse Transcriptase (Life Technologies, Carlsbad, CA, USA). For quantitative polymerase chain reaction (PCR), 100 ng of cDNA served as a template for PCR amplification using Brilliant SYBR green QPCR master mix according to the manufacturer's instructions (FastStart DNA Master SYBR Green I, Roche Molecular Biochemicals). Serial dilution (1 ng–1 fg/ μ L) of cDNA was used as a template for generation of a standard curve. Nested primers were used to amplify standards and kidney cDNA samples as follows: GGC TTC AAT TAC CCA CTG GA (sense) and TTG ATC CCA CAG CCA GTG TA (antisense) for AQP1 (498 bp)¹⁵⁾, TCC AGC AGT TGT

CAC TGG C (sense) and TAA GCA CAG TCC CCC AGA AGG (antisense) for AQP2 (562 bp)¹⁶⁾, CCA CAA TGG AAG GTT CTG TG (sense) and CCA ATC TCT AAG AAC CCT GT (antisense) for AQP7 (884 bp)¹⁷⁾, and TGC TGT GAC AGG TGT GTT AGG TCA (sense) and CCA TGG TTC TGC AAA TCG GG (antisense) for vasopressin-2 receptor (625 bp)¹⁸⁾.

Standards and unknown samples were amplified in duplicate in 96-well plates. The thermal profile of the LightCycler[®] Instrument (Roche Molecular Biochemicals, Mannheim, Germany) was optimized with an initial denaturation of 10 minutes at 95°C and 40 amplification cycles with each 15 seconds at 95°C, 5 seconds at 58°C and 10 seconds at 72°C. The comparative Ct method was used to determine the relative amounts of target-mRNA levels calculated for each sample by expressing the target-mRNA level as a percentage of GAPDH mRNA levels. Ct ratios were analyzed using the LightCycler[®] Software (Version 4.05). Specificity was ensured by postrun melting curve analysis.

6. Primary antibodies

For immunoblot analysis from rat kidneys, we used polyclonal antibodies against AQP1 and AQP2¹⁹⁾. They were kindly donated by Dr. Mark Knepper at the National Institutes of Health. In immunoblotting from cultured PTCs, commercially available antibodies (AQP1, Santa Cruz, cat. No. SC-20810; AQP7, Santa Cruz, cat. No. SC-28625) were used.

7. Statistics

Values are presented as means $\pm$ SE. Comparisons between cyclophosphamide-treated and control rats were performed by the Mann-Whitney U-test. *P* values of less than 0.05 were considered statistically significant.

Results

1. Effects of cyclophosphamide administration on rat kidneys

Table 1 shows blood and urine data obtained on the final day of Animal Experiment I. Compared with vehicle-treated controls, the 4 different doses (12, 24, 48, and 96 mg/kg BW) of cyclophosphamide produced no significant changes in urine output, urine osmolality, plasma sodium and creatinine concentration, and creatinine clearance.

The levels of AQP1 protein expression estimated in whole kidney homogenates were $100 \pm 18\%$, $136 \pm 53\%$, $244 \pm 38\%$, $614 \pm 194\%$, and $460 \pm 46\%$ at 0, 12, 24, 48, and 96 mg/kg BW doses, respectively. Thus, high doses of cyclophosphamide administration produced significant increases in AQP1 protein expression as compared with vehicle-treated controls ($P < 0.005$ at 48 mg/kg BW; $P < 0.05$ at 96 mg/kg BW). The expression of AQP2 protein also showed an increasing tendency at high doses of cyclophosphamide but did not reach the statistical significance (Fig. 1).

Table 2 shows blood and urine data obtained on the final day of Animal Experiment II. Although body weight

Table 1. Blood and Urine Data from Animal Experiment I

	0 mg/kg (n = 3)	12 mg/kg (n = 3)	24 mg/kg (n = 3)	48 mg/kg (n = 3)	96 mg/kg (n = 2)
Body weight (g)	259 $\pm$ 9	259 $\pm$ 4	251 $\pm$ 4	251 $\pm$ 5	247 $\pm$ 1
Urine volume (mL/100g BW)	15.8 $\pm$ 0.4	14.8 $\pm$ 0.2	18.8 $\pm$ 1.0	15.2 $\pm$ 2.1	10.5 $\pm$ 1.6
Urine sodium (μ mol/day/100g BW)	1060 $\pm$ 12	993 $\pm$ 81	1025 $\pm$ 166	994 $\pm$ 80	756 $\pm$ 31
Urine osmolality (mOsm/kg H ₂ O)	594 $\pm$ 25	544 $\pm$ 64	490 $\pm$ 43	628 $\pm$ 57	694 $\pm$ 112
Plasma sodium (mEq/L)	147.7 $\pm$ 0.3	147.3 $\pm$ 2.2	147.7 $\pm$ 0.7	148.3 $\pm$ 0.3	147.5 $\pm$ 0.5
Plasma creatinine (mg/dL)	0.57 $\pm$ 0.03	0.53 $\pm$ 0.03	0.53 $\pm$ 0.03	0.50 $\pm$ 0.00	0.50 $\pm$ 0.00
Creatinine clearance (μ L/min/g BW)	2.11 $\pm$ 0.09	1.94 $\pm$ 0.16	2.01 $\pm$ 0.12	1.85 $\pm$ 0.11	1.68 $\pm$ 0.01

Data are mean $\pm$ SE. No significant differences in all parameters were found among the groups when comparisons were done by Kruskal-Wallis test. BW, body weight.




Fig. 1. Immunoblot Analysis for Aquaporin-1 (AQP1) and Aquaporin-2 (AQP2) from Whole Kidneys of Animal Experiment I. Each lane was loaded with a protein sample from a different rat. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

* $P < 0.05$.

† $P < 0.005$ compared with vehicle-treated controls.

was steadily increased in both groups, weight gain was smaller in cyclophosphamide-treated rats than in controls ($P < 0.05$). Notably, cyclophosphamide-treated rats had lower plasma sodium concentration than controls (143 ± 1 vs. 146 ± 1 mEq/L, $P < 0.05$). However, urine output, urine osmolality and sodium concentration, plasma creatinine concentration, and creatinine clearance were not significantly different between the 2 groups.

Figure 2 shows the results of immunoblot analysis for




Fig. 2. Immunoblot Analysis for Aquaporin-1 (AQP1) and Aquaporin-2 (AQP2) from Renal Cortices of Animal Experiment II. Each lane was loaded with a protein sample from a different rat. No statistically significant differences were found between the groups.

Table 2. Blood and Urine Data from Animal Experiment II

Parameters	Control (n = 6)	Treated (n = 6)	P^*
Body weight (g)	306 ± 7	277 ± 4	< 0.05
Urine volume (mL/100g BW)	6.2 ± 1.0	4.3 ± 0.6	NS
Urine sodium ($\mu\text{mol/day/g BW}$)	8.0 ± 0.6	6.6 ± 0.6	NS
Urine osmolality (mOsm/kg H_2O)	830 ± 125	971 ± 108	NS
Plasma sodium (mEq/L)	146.3 ± 1.3	142.8 ± 0.8	< 0.05
Plasma creatinine (mg/dL)	0.63 ± 0.02	0.62 ± 0.02	NS
Creatinine clearance ($\mu\text{L/min/g BW}$)	2.36 ± 0.19	2.26 ± 0.26	NS

Data are mean $\pm$ SE. BW, body weight; NS, not significant.

*Comparisons by Mann-Whitney U test.

AQP1 and AQP2 from renal cortices of Animal Experiment II. Cyclophosphamide treatment caused insignificant increases in both AQP1 (136 ± 20 vs. $100 \pm 23\%$) and AQP2 (157 ± 18 vs. $100 \pm 22\%$) protein expression. Figure 3 illustrates the results of reverse transcription PCR for AQP1, AQP2, vasopressin-2 receptor, and aquaporin-7 (AQP7) from each RNA sample of renal cortex. QPCR,




Fig. 3. Reverse Transcription Polymerase Chain Reaction (PCR) from Renal Cortices of Animal Experiment II. Each lane was loaded with a cDNA sample from a different rat. AQP1, aquaporin-1; AQP2, aquaporin-2; V₂R, vasopressin-2 receptor; AQP7, aquaporin-7; GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

however, revealed no significant differences in mRNA levels between treated versus control rats (Fig. 4): AQP1, 75 ± 11 vs. $100 \pm 21\%$; AQP7, 105 ± 16 vs. $100 \pm 13\%$; AQP2, 81 ± 14 vs. $100 \pm 13\%$; and vasopressin-2 receptor, 96 ± 33 vs. $100 \pm 13\%$.

2. Effects of 4-HC treatment on rabbit PTCs

In order to examine the effect of an active metabolite of cyclophosphamide on the expression of water channel proteins in primary cultured renal PTCs, we treated rabbit PTCs with 4-HC ($0\text{--}30 \mu\text{mol/L}$) for 24 hours. In immunoblotting using membrane fraction of PTCs, protein expression levels of AQP1 and AQP7 were increased in a dose-dependent manner by 4-HC treatment (Fig. 5).

Discussion

The kidney is able to regulate water excretion to maintain water balance. Renal conservation of water occurs partly in association with sodium reabsorption in the proximal tubule and thin limb of Henle's loop and partly as a result of action of vasopressin on the collecting duct. AQP1 and AQP2 are the major water channels to mediate water reabsorption in the proximal nephron and collecting duct,




Fig. 4. Quantitative Polymerase Chain Reaction (PCR) from Renal Cortices of Animal Experiment II. AQP1, aquaporin-1; AQP2, aquaporin-2; AQP7, aquaporin-7; V₂R, vasopressin-2 receptor. No statistically significant differences were found between the groups.

respectively.

Excessive accumulation of water results in dilutional hyponatremia. It usually occurs due to impaired renal excretion of water rather than excessive water intake because the capacity for water excretion normally is so great. When drugs affect water homeostasis, they can impair the renal diluting capacity in 3 ways²⁰⁾. The first possibility




Fig. 5. Immunoblot for Aquaporin-1 (AQP1) and Aquaporin-7 (AQP7) from Primary Cultured Rabbit Renal Proximal Tubule Cells. Cells were treated for 24 hours with each concentration of 4-hydroperoxycyclophosphamide (4-HC).

is that they can increase vasopressin secretion centrally. However, plasma vasopressin levels are not elevated in patients following the administration of intravenous cyclophosphamide^{21,22}. In addition, antidiuresis was reported to occur in response to intravenous cyclophosphamide in an 8 year-old girl with central diabetes insipidus²³, excluding the possibility of a drug-induced syndrome of inappropriate antidiuretic hormone secretion (SIADH).

The second possibility is reset osmostat, lowering the threshold for vasopressin secretion. This condition is usually associated with a long-term course of hyponatremia such as spinal cord injury, psychiatric problems, and alcoholism²⁴, but cyclophosphamide-induced water retention is an acute process. The injection of cyclophosphamide into normally hydrated patients caused antidiuresis beginning in 4 to 12 hours and lasting as long as 20 hours²⁵.

The third possibility is that cyclophosphamide may potentiate the action of vasopressin on the kidney. This postulation is very plausible, but has never been verified for decades. Recently, only carbamazepine was reported to increase AQP2 expression²⁶. In this study, the protein expression of AQP2 seemed to increase, although statistical significance was not reached. Further studies will be required to demonstrate the role of AQP2 in cyclophosphamide-induced hyponatremia.

This study was focused to elucidate the role of the AQP1 water channel in cyclophosphamide-induced water retention. We tested whether AQP1 protein may be affected by cyclophosphamide administration in rat kidneys. In addition, primary cultured renal PTCs were treated with 4-HC because the cyclophosphamide-induced antidiuresis was related to the urinary excretion of active alkylating

metabolites of the drug²⁵.

To avoid the occurrence of hemorrhagic cystitis, we used moderate doses of a single cyclophosphamide administration. In Animal Experiment I, the increase in AQP1 protein expression was remarkable when 48 and 96 mg/kg BW cyclophosphamide were administered (Fig. 1). However, hyponatremia was not found in these rats (Table 1). Considering that cyclophosphamide-induced hyponatremia usually occurs 4–12 hours after the intravenous administration of the drug, we may have to test an earlier change in plasma sodium level.

The results of Animal Experiment II suggest that the increase in AQP1 protein may persist for more than several days after a single cyclophosphamide administration (Fig. 2). Cyclophosphamide-treated rats had a significantly lower plasma sodium concentration. Although weight gain was smaller in these animals, urine sodium excretion was not different between treated and control rats (Table 1). Thus we think that the main cause of hyponatremia in these cyclophosphamide-treated rats is water retention rather than hypovolemia.

Consistent with the animal studies, we found that treatment of primary cultured renal PTCs with an active metabolite of cyclophosphamide (4-HC) increased protein expressions of AQP1 and AQP7 in a dose-dependent manner (Fig. 5). Previous studies reported that AQP7 was localized selectively at the brush border membranes of proximal straight tubule (segment 3 proximal tubule, S3) in the rat kidney, suggesting that AQP7 may function as a pathway for transcellular water transport in S3 in concert with more widely expressed AQP1 in proximal tubules^{27,28}. Thus, the proximal nephron should be one of the major target sites where cyclophosphamide-induced water retention occurs.

Despite the notable changes in AQP1 and AQP7 protein, we found no differences in the mRNA levels between cyclophosphamide-treated and control rats (Fig. 4). It is conceivable that other mechanisms such as decreased protein degradation than transcriptional regulation may exert to affect the AQP1 protein level. Further studies

are required to demonstrate intracellular mechanisms to increase the protein abundance of AQP1 in response to cyclophosphamide administration.

Acknowledgements

This study was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (2010-0011799).

References

- 1) Moses AM, Miller M: Drug-induced dilutional hyponatremia. *N Engl J Med* 291:1234-1239, 1974
- 2) Steele TH, Serpick AA, Block JB: Antidiuretic response to cyclophosphamide in man. *J Pharmacol Exp Ther* 185:245-253, 1973
- 3) Koo TY, Bae SC, Park JS, et al.: Water intoxication following low-dose intravenous cyclophosphamide. *Electrolyte Blood Press* 5:50-54, 2007
- 4) Lee YC, Park JS, Lee CH, et al.: Hyponatraemia induced by low-dose intravenous pulse cyclophosphamide. *Nephrol Dial Transplant* 25:1520-1524, 2010
- 5) Nielsen S, Agre P: The aquaporin family of water channels in kidney. *Kidney Int* 48:1057-1068, 1995
- 6) Nielsen S, DiGiovanni SR, Christensen EI, Knepper MA, Harris HW: Cellular and subcellular immunolocalization of vasopressin-regulated water channel in rat kidney. *Proc Natl Acad Sci U S A* 90:11663-11667, 1993
- 7) Lee YJ, Han HJ: Troglitazone ameliorates high glucose-induced EMT and dysfunction of SGLTs through PI3K/Akt, GSK-3 β , Snail1, and β -catenin in renal proximal tubule cells. *Am J Physiol Renal Physiol* 298:F1263-1275, 2010
- 8) Bodo E, Tobin DJ, Kamenisch Y, et al.: Dissecting the impact of chemotherapy on the human hair follicle: a pragmatic in vitro assay for studying the pathogenesis and potential management of hair follicle dystrophy. *Am J Pathol* 171:1153-1167, 2007
- 9) Flowers JL, Ludeman SM, Gamcsik MP, et al.: Evidence for a role of chloroethylaziridine in the cytotoxicity of cyclophosphamide. *Cancer Chemother Pharmacol* 45:335-344, 2000
- 10) Anderson LW, Ludeman SM, Colvin OM, Grochow LB, Strong JM: Quantitation of 4-hydroxycyclophosphamide/aldophosphamide in whole blood. *J Chromatogr B Biomed Appl* 667:247-257, 1995
- 11) Anderson LW, Chen TL, Colvin OM, et al.: Cyclophosphamide and 4-Hydroxycyclophosphamide/aldophosphamide kinetics in patients receiving high-dose cyclophosphamide chemotherapy. *Clin Cancer Res* 2:1481-1487, 1996
- 12) Chen TL, Kennedy MJ, Anderson LW, et al.: Nonlinear pharmacokinetics of cyclophosphamide and 4-hydroxycyclophosphamide/aldophosphamide in patients with metastatic breast cancer receiving high-dose chemotherapy followed by autologous bone marrow transplantation. *Drug Metab Dispos* 25:544-551, 1997
- 13) Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-254, 1976
- 14) Lee CH, Kim S, Kang CM, Kim WY, Kim J, Kim GH: Altered expression of tight junction proteins in cyclosporine nephrotoxicity. *Am J Nephrol* 33:7-16, 2011
- 15) Sonoda H, Yokota-Ikeda N, Oshikawa S, et al.: Decreased abundance of urinary exosomal aquaporin-1 in renal ischemia-reperfusion injury. *Am J Physiol Renal Physiol* 297:F1006-1016, 2009
- 16) Kishore BK, Mandon B, Oza NB, et al.: Rat renal arcade segment expresses vasopressin-regulated water channel and vasopressin V2 receptor. *J Clin Invest* 97:2763-2771, 1996
- 17) Nejsum LN, Elkjaer M, Hager H, Frokiaer J, Kwon TH, Nielsen S: Localization of aquaporin-7 in rat and mouse kidney using RT-PCR, immunoblotting, and immunocytochemistry. *Biochem Biophys Res Commun* 277:164-170, 2000
- 18) Sarmiento JM, Ehrenfeld P, Anazco CC, et al.: Differential distribution of the vasopressin V receptor along the rat nephron during renal ontogeny and maturation. *Kidney Int* 68:487-496, 2005
- 19) Terris J, Ecelbarger CA, Nielsen S, Knepper MA: Long-term regulation of four renal aquaporins in rats. *Am J Physiol* 271:F414-422, 1996
- 20) Liamis G, Milionis H, Elisaf M: A review of drug-induced hyponatremia. *Am J Kidney Dis* 52:144-153, 2008
- 21) Bode U, Seif SM, Levine AS: Studies on the antidiuretic effect of cyclophosphamide: vasopressin release and sodium excretion. *Med Pediatr Oncol* 8:295-303, 1980
- 22) Bressler RB, Huston DP: Water intoxication following moderate-dose intravenous cyclophosphamide. *Arch Intern Med* 145:548-549, 1985
- 23) Campbell DM, Atkinson A, Gillis D, Sochett EB: Cyclophosphamide and water retention: mechanism revisited. *J Pediatr Endocrinol Metab* 13:673-675, 2000
- 24) Kahn T: Reset osmostat and salt and water retention in the course of severe hyponatremia. *Medicine (Baltimore)* 82:170-176, 2003

- 25) DeFronzo RA, Braine H, Colvin M, Davis PJ: Water intoxication in man after cyclophosphamide therapy. Time course and relation to drug activation. *Ann Intern Med* 78:861-869, 1973
- 26) de Bragança AC, Moyses ZP, Magaldi AJ: Carbamazepine can induce kidney water absorption by increasing aquaporin 2 expression. *Nephrol Dial Transplant* 25:3840-3845, 2010
- 27) Ishibashi K, Imai M, Sasaki S: Cellular localization of aquaporin 7 in the rat kidney. *Exp Nephrol* 8:252-257, 2000
- 28) Nejsum LN, Elkjaer M, Hager H, Frokiaer J, Kwon TH, Nielsen S: Localization of aquaporin-7 in rat and mouse kidney using RT-PCR, immunoblotting, and immunocytochemistry. *Biochem Biophys Res Commun* 277:164-170, 2000