

Severe Hyponatremia by Excessive Bamboo Salt Ingestion in Healthy Young Woman

Hyun Jun Ju, M.D., Hong Jin Bae, M.D.,
Dae Eun Choi, M.D., Ki Ryang Na, M.D.,
Kang Wook Lee, M.D., Young Tai Shin, M.D.
Renal Division, Department of Internal Medicine,
Chungnam National University Hospital,
Daejeon, Korea

Received: January 1, 2013

Accepted: October 18, 2013

Corresponding Author: Kang Wook Lee, M.D.
Department of Internal Medicine, Chungnam
National University Hospital, 640 Daesa-Dong,
Chung-Ku, Daejeon 301-721, Korea
Tel: +82-41-412-6739, Fax: +82-41-412-5574
E-mail: kwlee@cnu.ac.kr

Severe hyponatremia is an important electrolyte disorder that has serious effects. The patient had no medical history. A 20-year-old ingested bamboo salt for digestion and weight reduction according to the folk remedies posted on an internet website. She presented with vomiting and diarrhea over ten times per day. Her initial serum sodium concentration was 174 mEq/L. Her symptoms improved rapidly with hypotonic saline infusion. She recovered completely without any sequelae in three days. Severe hyponatremia in a normal young adult with clear consciousness and normal renal function has not been reported in Korea yet. So we report a case of severe hyponatremia by excessive bamboo salt ingestion in healthy young woman.

Key Words: Hyponatremia, Folk remedies

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Most of the known hyponatremic cases are due to a loss of water rather than due to excessive salt intake. The toxicity of large amounts of salt have been known for a long time. It is reported that in ancient China, saturated salt solutions were used as a traditional suicide method. The causes of high salt intake induced hyponatremia were gastric lavage with high salt solution, excessive salt intake, intravenous infusion of hypertonic saline, forced salt intake in cases of child abuse or exorcism¹⁾ and accidental sodium chloride ingestion due to inadvertent preparation of child food with salt instead of sugar²⁾. Recently, a case of severe hyponatremia caused by excessive salt intake as folk remedies for 3 months in a patient with chronic kidney disease had been reported in Korea³⁾. However, severe hyponatremia in a healthy young adult with clear consciousness and normal renal function had not been

reported in Korea yet. We report a case of a healthy young woman with acute severe hyponatremia caused by excessive bamboo salt intakes as folk remedies.

Case Report

The patient was a 20-year-old previously healthy woman. She was a university student. Her height and body weight were 168.1 cm and 52.2 kg, respectively. She ingested bamboo salt (about 150 grams) in a day for the purpose of digestion and weight reduction. After that, she had more than ten episodes of vomiting and diarrhea for 2-3 hours. On arrival at our emergency department, vital signs were blood pressure of 128/81 mmHg, heart rate of 75 beat/min, respiration rate of 20/min, and body temperature of 37.3°C. Her consciousness was clear but she complained of generalized weakness. The complete blood counts showed hemoglobin 11.3 g/dL, WBC 10,000 cells/ μ L, and platelet 196,000/ μ L. The results of her blood

chemistry were as follows: Aspartate transaminase 11 IU/L, Alanine transaminase 9 IU/L, total protein 6.4 g/dL, albumin 4.5 g/dL, blood urea nitrogen 5.2 mg/dL, and creatinine 0.57 mg/dL. Serum osmolarity was 348 mosm/kg. Serum sodium, potassium and chloride were 174 mEq/L, 3.6 mEq/L and 135 mEq/L respectively. Urine sodium, potassium, chloride and osmolarity were 440 mEq/L, 125.7 mEq/L, 314 mEq/L and 841 mosm/kg, respectively. On arterial blood gas analysis, pH was 7.32, PaCO₂ 36 mmHg, bicarbonate 18.5 mEq/L, PaO₂ 93 mmHg and O₂ saturation 97%. On the 1st day, total input and urine output was 3,700 ml and 2,100 ml, respectively. Her symptoms such as nausea, vomiting and diarrhea improved rapidly with 0.45% saline infusion and supportive care. On the 2nd day, her serum sodium concentration dropped to 152 mEq/L and urine output was 2,200 ml/day. On the 3rd day, serum sodium concentration returned to normal range (Fig 1). She recovered completely without any sequelae in 3 days.

Discussion

Hyponatremia is characterized by a decrease in total body water relative to total body electrolyte contents resulting in the increase of sodium concentration of extracellular fluids. Recently, hyponatremia resulting from excessive sodium intake have been described after using salt as folk remedies for three months in a patient with chronic

kidney disease in Korea³.

In Korean folk remedies, bamboo salt was believed to have many beneficial effects on digestion, inflammation, detoxification and so on. Sohn et al. showed that bamboo salt had anti-plaque and anti-inflammatory effect on oral hygiene used as dentifrices⁴. Sea salt is stuffed into bamboo tubes, and the ends are plugged with mud; the assembly is roasted one or more times to make bamboo salt. So the main ingredient of bamboo salt is sodium chloride salt. And it is believed that the trace elements in the mud and bamboo are thought to make this form of salt healthier. However Yoo et al. reported that bamboo salt had no effects on the general pharmacology of central nervous systems and cardiovascular system⁵. So it is probably reasonable that bamboo salt maybe regarded as sodium chloride salt in the pharmacologic aspect.

The mortality of hyponatremia is very high. In general, doses greater than 1 g sodium chloride per kg body weight can lead to raised blood sodium concentration by 30 mmol/L⁶⁻⁸. Serum sodium levels above 160 mEq/L have been associated with a high mortality rate (9-35%) and neurological damage⁹⁻¹¹. Moreover, extreme hyponatremia (>190 mEq/L) has a high mortality rate up to 62% for children¹².

Severe hyponatremia by excessive salt ingestion occurs acutely and have high mortality rate. Paut et al. report a survivor from extreme hyponatremia in accidental salt poisoning in an infant¹³. Ofran et al. described highest sodium plasma level of 255 mEq/L in a female with psychiatric disorder⁷.

Acute hyponatremia should be treated with rapid rate of serum concentration of 1-2 mEq/L/h, whereas chronic hyponatremia should be treated with a slower rate of sodium concentration of 0.5 mEq/L/h. However, the over-correction of hyponatremia might cause severe complications, because some acute hyponatremia can not be differentiated from chronic hyponatremia¹⁴. We administered hypotonic saline for just 2 days. The serum sodium concentration on admission day (174 mEq/L) decreased to 152 mEq/L within 1 day. But she recovered completely without any sequelae. We assume that her young age and healthy condition contributed to this favorable result¹⁵.




Fig. 1. Serial data of serum sodium level and osmolarity during clinical course. sOsm; serum osmolarity, sNa; serum sodium

In summary, we present here a case of severe hypernatremia by excessive bamboo salt ingestion in a healthy young woman with clear consciousness and completely recovered without any sequelae.

References

1. Hedouin V, Revuelta E, Becart A, Tournel G, Deveaux M, Gosset D: A case of fatal salt water intoxication following an exorcism session. *Forensic Sci Int* 99(1):1-4, 1999
2. Johnston JG, Robertson WO: Fatal ingestion of table salt by an adult. *West J Med* 126(2):141-143, 1977
3. Jang JS, Choi HM, Hyun YY, Kwon BS, Park JY, Kim JH, Lee JW, Lee JW, Kwon YJ, Pyo HJ: A case of Hypernatremia by folk remedies in a CRF patient. *Korean J Nephrol* 25(4):675-679, 2006
4. Sohn WS, Chung SC, Kim CY. Comparison of the anti-plaque and anti-inflammatory effect of the dentifrices containing NaCl and bamboo salt. *J Kor Dental Health* 15(2), 000-000, 1991
5. Yoo TM, Kim SS, Roh YN, Yi SY, Kim OK, Rheu HM, Yang JS: General pharmacology of bamboo salt. *J Appl Pharmacol* 8:93-98, 2000
6. Turk EE, Schulz F, Koops E, Gehl A, Tsokos M. Fatal hypernatremia after using salt as an emetic - - report of three autopsy cases. *Leg Med (Tokyo)* 7(1):47-50 2005
7. Ofra Y, Lavi D, Opher D, Weiss TA, Elinav E: Fatal voluntary salt intake resulting in the highest ever documented sodium plasma level in adults (255 mmol L⁻¹): a disorder linked to female gender and psychiatric disorders. *J Intern Med* 256(6):525-528, 2004
8. Ward DJ: Fatal hypernatremia after a saline emetic. *Br Med J* 2(5354):432, 1963
9. Snyder NA, Feigal DW, Arief AI: Hypernatremia in elderly patients. A heterogeneous, morbid, and iatrogenic entity. *Ann Intern Med* 107(3):309-319, 1987
10. Meadow R: Non-accidental salt poisoning. *Arch Dis Child* 68(4):448-452, 1993
11. Dine MS, McGovern ME: Intentional poisoning of children an overlooked category of child abuse: report of seven cases and review of the literature. *Pediatrics* 70(1): 32-35, 1982
12. Moder KG, Hurley DL: Fatal hypernatremia from exogenous salt intake: report of a case and review of the literature. *Mayo Clin Proc* 65(12):1587-1594, 1990
13. Paut O, Andre N, Fabre P, Sobraques P, Drouet G, Arditti J, Camboulives J: The management of extreme hypernatraemia secondary to salt poisoning in an infant. *Paediatr Anaesth* 9(2):171-174, 1999
14. Star RA. Hyperosmolar states. *Am J Med Sci* 300(6):402-412, 1990
15. Casavant MJ, Fitch JA: Fatal hypernatremia from salt-water used as an emetic. *J Toxicol Clin Toxicol* 41(6): 861-863, 2003