

Hypertension Management in Patients with Chronic Kidney Disease in the Post-SPRINT Era

Hae Hyuk Jung

Department of Medicine, Kangwon National University Hospital, Kangwon National University School of Medicine, Chuncheon, Republic of Korea

Received: October 2, 2021

Revised: October 27, 2021

Accepted: November 15, 2021

Corresponding Author: Hae Hyuk Jung, MD, PhD
Department of Medicine, Kangwon National University Hospital, 156 Baekryung-ro, Chuncheon, Gangwon-do 24289, Korea
Tel: +82-33-258-2376; Fax: +82-33-258-2455
E-mail: haehyuk@kangwon.ac.kr

The management of high blood pressure (BP) is crucial for improving outcomes in patients with chronic kidney disease (CKD). The updated Kidney Disease: Improving Global Outcomes 2021 BP guideline proposes treating adults with CKD to a target systolic BP (SBP) of <120 mmHg based on the standardized office BP measurement. This suggestion is largely based on the finding of SPRINT (Systolic Blood Pressure Intervention Trial) that targeting an SBP of <120 mmHg versus <140 mmHg is beneficial for cardiovascular and mortality outcomes, regardless of the patient's kidney disease status. However, extended follow-up studies of CKD trials showed that intensive versus usual BP control was associated with a lower risk of kidney failure in patients with, but not in those without, proteinuria. Similarly, a recent population-based study in Korea demonstrated that the optimal on-treatment BP for composite cardiorenal and mortality outcomes was left-shifted in adults with CKD, particularly in those with albuminuria, relative to that in patients without CKD. Moreover, in meta-analyses of randomized trials, more intensive versus standard BP control was associated with a lower risk of all-cause mortality in patients with CKD and albuminuria but not in those without CKD. Meanwhile, a 2020 Cochrane review reported that lower BP targets ($\leq 135/85$ mmHg), compared with standard targets ($\leq 140/90$ mmHg), resulted in a small reduction in cardiovascular events, an increase in other serious adverse events, and no reduction in total serious adverse events. Lowering SBP to <120 mmHg can potentially increase the risk of treatment-related adverse events beyond the cardioprotective benefits, and standardized BP measurement increases the burden on patients and resources. Thus, targeting a BP of <130/80 mmHg with appropriate office BP measurement can be an option in patients with CKD. The presence of albuminuria would need to be additionally considered to determine individualized BP targets.

Key Words: Antihypertensive agents, Blood pressure, Chronic kidney disease, Practice guideline

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Chronic kidney disease (CKD) is a common health problem worldwide and is associated with a high risk of adverse renal and cardiovascular events. Hypertension is common in patients with CKD, and the management of blood pressure (BP) is known to be crucial for slowing the progression of kidney disease and for reducing the rates of cardiovascular events. In early 2021, the Kidney Disease: Improving Global Outcomes (KDIGO) released an updated BP guideline that

suggests a target systolic BP (SBP) of <120 mmHg based on the standardized office measurement in adults with CKD¹. However, this suggestion is largely based on the results of a single trial, SPRINT (Systolic Blood Pressure Intervention Trial), and standardized BP measurements require efforts from patients and health-care providers. The limitations of the evidence and gaps between guidelines and practice will be reviewed here.

Table 1. Extended follow-up studies for intensive versus usual BP control in CKD

| Clinical Trial (Follow-up) | Target MAP, mmHg | Baseline Kidney Function | Progression to ESKD, HR (95% CI) | All-cause Death, HR (95% CI) |
|--|-----------------------|--|----------------------------------|------------------------------|
| CKD with Severe Proteinuria | | | | |
| MDRD-a subset ³⁾ (median 10.7 y) | <92 vs. <107 | Urine protein ≥ 1 g/d and GFR 13-55 | 0.55 (0.41-0.72) | NA |
| AASK subset ⁶⁾ (median 14.4 y) | ≤ 92 vs. 102-107 | Urine protein ≥ 1 g/d and GFR 20-65 | 0.59 (0.41-0.85) | NA |
| CKD without Severe Proteinuria | | | | |
| MDRD-a subset ³⁾ (median 10.7 y) | <92 vs. <107 | Urine protein <1 g/d and GFR 13-55 | 0.81 (0.64-1.02) | NA |
| AASK subset ⁶⁾ (median 14.4 y) | ≤ 92 vs. 102-107 | Urine protein <1 g/d and GFR 20-65 | 1.05 (0.83-1.32) | NA |
| CKD, Total Participants | | | | |
| MDRD-b ⁵⁾ (median 19.3 y) | <92 vs. <107 | GFR 13-55, urine protein ≥ 0.3 g/d in 51.8% | 0.86 (0.73-1.00) | 0.82 (0.68-0.98) |
| AASK ⁶⁾ (median 14.4 y) | ≤ 92 vs. 102-107 | GFR 20-65, PCR ≥ 0.22 in 32.6% | 0.92 (0.75-1.12) | 0.92 (0.77-1.10) |

*Data are for all participants with or without severe proteinuria.

BP, blood pressure; CI, confidence interval; ESKD, end-stage kidney disease; GFR, glomerular filtration rate (mL/min/1.73 m²); HR, hazardratio; MAP, mean arterial pressure; NA, not available; PCR, urineprotein-to-creatinineratio (g/g); s-Cr, serum creatinine.

The KDIGO 2021 BP Guideline

The KDIGO BP guideline recommends starting renin-angiotensin system inhibitors (RASi) (angiotensin-converting enzyme inhibitors [ACEi] or angiotensin II receptor blockers [ARB]) in persons with high BP, CKD, and severely increased albuminuria (estimated glomerular filtration rate [eGFR] categories G1-G4, urine albumin-to-creatinine ratio [ACR] category A3) without diabetes and in persons with high BP, CKD, and moderately to severely increased albuminuria (G1-G4, A2-A3) with diabetes. The guideline suggests starting ACEi or ARB in persons with high BP, CKD, and moderately increased albuminuria (G1-G4, A2) without diabetes. Furthermore, the guideline recommends avoiding any combination of ACEi, ARB, and direct renin inhibitor therapy in persons with CKD, with or without diabetes.

The key points of the 2021 guideline are the recommendation of a standardized office BP measurement instead of a routine office BP measurement for the management of high BP in adults and the suggestion of treating adults with high BP and CKD to a target SBP of <120 mmHg,

when tolerated, based on the standardized office BP measurement. The guideline suggests using out-of-office BP measurements with ambulatory BP monitoring or home BP monitoring as complementary readings to standardized office BP readings.

In comparison, the original KDIGO BP guideline recommends maintaining an office BP of <130/80 mmHg in patients with CKD and albuminuria and <140/90 mmHg in patients with CKD without albuminuria²⁾. This recommendation was based on the results of extended follow-up studies of randomized trials in patients with CKD (Table 1)³⁻⁶⁾. Conversely, the recommendation of a target SBP of <120 mmHg with the standardized measurement is largely based on the findings of SPRINT in CKD subsets, in which BP was determined as an average of 3 readings obtained after a rest period of 5 min, with the patients sitting with their back supported and without talking⁷⁾. Standardized BP measurements increase the burden on patients, health-care providers, and facilities. Although ambulatory or home BP monitoring can be used to complement standardized office BP measurements, no completed large clinical trials have recommended an out-of-office BP target.

Table 2. Renin-angiotensin system inhibitors in CKD

| Clinical Trial (Index Disease) | Treatment vs. Control Group | Baseline Kidney Function | CKD Progression, Risk Reduction % | All-cause Death, Rate % |
|--|--------------------------------------|--|---------------------------------------|-------------------------|
| Benazepril ⁸⁾ (nondiabetic CKD) | Benazepril vs. placebo | eGFR 30-60, mean urine protein 1.8 g/d | 53% for s-Cr doubling or ESKD | 2.7% vs. 0.4% |
| Benazepril ¹²⁾ (nondiabetic CKD) | Benazepril vs. placebo | s-Cr 3.1-5.0 mg/dL and urine protein \geq 0.3 g/d | 48% for s-Cr doubling or ESKD | 0.9% vs. 0.0% |
| REIN ⁹⁾ (nondiabetic CKD) | Ramipril vs. placebo | Urine protein 1.0-3.0 g/d and CrCl 20-70 | 56% for ESKD | 1.0% vs. 0.0% |
| REIN ¹⁰⁾ (nondiabetic CKD) | Ramipril vs. placebo | Urine protein \geq 3.0 g/d and CrCl 20-70 | 49% for s-Cr doubling or ESKD | 2.6% vs. 1.1% |
| AASK ¹⁶⁾ (hypertensive CKD) | Ramipril vs. amlodipine | GFR 20-65, PCR >0.22 in 32.6% | 38% for GFR decline or ESKD | 1.5% vs. 1.7% |
| Captopril ¹³⁾ (DKD, T1D) | Captopril vs. placebo | Urine protein \geq 0.5 g/d and s-Cr \leq 2.5 mg/dL | 50% for ESKD or death | 3.9% vs. 6.9% |
| RENAAL ¹⁴⁾ (DKD, T2D) | Losartan vs. placebo | ACR \geq 0.3 and s-Cr 1.0-3.0 mg/dL | 21% for s-Cr doubling or ESKD | 21.0% vs. 20.3% |
| IDNT ¹⁵⁾ (DKD, T2D) | Irbesartan vs. amlodipine or placebo | Urine protein \geq 0.9 g/d and s-Cr 1.0-3.0 mg/dL | 21% for s-Cr doubling, ESKD, or death | 15.0% vs. 15.5% |

ACEi, angiotensin-converting enzyme inhibitor; ACR, urine albumin-to-creatinine ratio (g/g); ARB, angiotensin receptor blocker; CKD, chronic kidney disease; CrCl, creatinine clearance (mL/min/1.73 m²); DKD, diabetic kidney disease; eGFR, estimated glomerular filtration rate (mL/min/1.73 m²); ESKD, end-stage kidney disease; GFR, glomerular filtration rate (mL/min/1.73 m²); PCR, urine protein-to-creatinine ratio (g/g); s-Cr, serum creatinine; T1D, type 1 diabetes; T2D, type 2 diabetes.

Renin-Angiotensin-Aldosterone System Inhibitors

Randomized controlled trials have shown the clinical benefits of RASi in patients with CKD and proteinuria (Table 2). The renoprotective effect of ACEi was demonstrated in trials for nondiabetic kidney disease with overt proteinuria⁸⁻¹⁰⁾, and the benefit persisted even in patients with advanced CKD (eGFR <30 mL/min/1.73 m²)^{11,12)}. The beneficial effect of ACEi was also shown in a trial for type 1 diabetes with proteinuria (urine protein \geq 500 mg/day)¹³⁾. The renoprotective effect of ARB was demonstrated in trials for type 2 diabetes with severely increased albuminuria (urine ACR \geq 300 mg/g)^{14,15)}.

However, combination therapy with ACEi and ARB increased the risks of adverse events, such as acute kidney injury and severe hyperkalemia, without providing clinical benefits in patients at a high risk of vascular events and in those with type 2 diabetes and severely increased albuminuria (Table 3)^{17,18)}. Similarly, treatment with aliskiren, a direct renin inhibitor, in combination with either an ACEi

or ARB resulted in more adverse events with no additional benefits in trials for type 2 diabetes with CKD or cardiovascular disease and for heart failure with an ejection fraction of \leq 35%^{19,20)}. In contrast, finerenone, a selective nonsteroidal mineralocorticoid receptor antagonist, in combination with an ACEi or ARB showed benefits in terms of slowing the progression of kidney disease and reducing the rates of cardiovascular events in trials for type 2 diabetes with CKD^{21,22)}.

New Classes of Drugs

Besides RASi, other new classes of drugs have shown a noticeable renoprotective benefit in randomized controlled trials (Table 4). Sacubitril-valsartan, an angiotensin receptor-neprilysin inhibitor (ARNi), reduced the risk of kidney disease progression and hyperkalemia compared with either an ACEi or ARB in patients with heart failure^{23,24)}. ARNi can also be considered in patients with CKD at a risk of hyperkalemia, although direct evidence in CKD would be needed to confirm the benefit.

Table 3. Combination therapy with renin-angiotensin-aldosterone system inhibitors

| Clinical Trial (Index Disease) | Treatment vs. Control Group | Participants | Kidney Disease Progression, HR (95% CI) | All-cause Death, HR (95% CI) | Hyperkalemia,* Rate % |
|--|--|---|--|------------------------------|--|
| ONTARGET ¹⁷⁾ (high risk patients) | Ramipril-telmisartan vs. ramipril | Vascular disease or high risk of diabetes | 1.24 (1.01-1.51) for s-Cr doubling or all dialysis | 1.07 (0.98-1.16) | 5.6% vs. 3.3% for s-K ⁺ >5.5 mmol/L |
| VA NEPHRON-D ¹⁸⁾ (DKD, T2D) | Losartan-lisinopril vs. losartan-placebo | ACR ≥0.3 and eGFR 30-90 | 0.78 (0.58-1.05) for eGFR decline or ESKD | 1.04 (0.73-1.49) | 9.9% vs. 4.4% |
| ALTITUDE ¹⁹⁾ (DKD, T2D) | Aliskiren vs. placebo [†] | High risk of cardiovascular and renal events | 1.03 (0.87-1.23) for s-Cr doubling or ESKD | 1.06 (0.92-1.23) | 11.2% vs. 7.2% |
| ATMOSPHERE ²⁰⁾ (heart failure) | Aliskiren-enalapril vs. enalapril | Chronic heart failure with ejection fraction ≤35% | 1.50 (0.82-2.74) for s-Cr doubling, ESKD, or renal death | 0.91 (0.82-1.02) | 5.0% vs. 3.6% |
| FIDELIO-DKD ²¹⁾ (DKD, T2D) | Finerenone vs. placebo [†] | “ACR 0.03-0.3, eGFR 25-60, and retinopathy” or “ACR 0.3-5 and eGFR 25-75” | 0.82 (0.73-0.93) for eGFR decline, ESKD, or renal death | 0.90 (0.75-1.07) | 4.5% vs. 1.4% |
| FIGARO-DKD ²²⁾ (DKD, T2D) | Finerenone vs. placebo [†] | “ACR 0.03-0.3 and eGFR 25-90” or “ACR 0.3-5 and eGFR 25-75” | 0.87 (0.76-1.01) for eGFR decline, ESKD, or renal death | 0.89 (0.77-1.04) | 2.3% vs. 1.2% |

*Data are rates for s-K⁺ >6.0 mmol/L unless otherwise stated.

[†]All patients were treated with either an angiotensin-converting enzyme inhibitor or an angiotensin II receptor blocker.

ACR, urine albumin-to-creatinine ratio (g/g); CI, confidence interval; DKD, diabetic kidney disease; eGFR, estimated glomerular filtration rate (mL/min/1.73 m²); ESKD, end-stage kidney disease; GFR, glomerular filtration rate; HR, hazard ratio; s-Cr, serum creatinine; s-K⁺, serum potassium; T2D, type 2 diabetes.

Table 4. Angiotensin receptor-neprilysin inhibitors in heart failure and sodium-glucose transporter-2 inhibitors in CKD

| Clinical Trial (Index Disease) | Treatment vs. Control Group | Participants | Kidney Disease Progression, HR (95% CI) | All-cause Death, HR (95% CI) | Other Outcomes, Rate % [*] |
|--|------------------------------------|--|--|------------------------------|--|
| PARAGON-HF ²³⁾ (heart failure) | Sacubitril-valsartan vs. valsartan | Symptoms of heart failure and ejection fraction ≥45% | 0.50 (0.33-0.77) for eGFR decline, ESKD, or renal death | 0.97 (0.84-1.13) | 3.1% vs. 4.3% for s-K ⁺ >6.0 mmol/L |
| PARADIGM-HF ²⁴⁾ (heart failure) | Sacubitril-valsartan vs. enalapril | Symptoms of heart failure and ejection fraction ≤40% | 0.86 (0.65-1.13) for eGFR decline or ESKD | 0.84 (0.76-0.93) | 4.3% vs. 5.6% for s-K ⁺ >6.0 mmol/L |
| CREDESCENCE ²⁹⁾ (DKD, T2D) | Canagliflozin vs. placebo | ACR 0.3-5 and eGFR 30-90 | 0.66 (0.53-0.81) for s-Cr doubling, ESKD, or renal death | 0.83 (0.68-1.02) | 33.5% vs. 36.7% for total serious adverse events |
| DAPA-CKD ³¹⁾ (CKD with albuminuria) | Dapagliflozin vs. placebo | ACR 0.2-5 and eGFR 25-75, T2D in 67.5% | 0.56 (0.45-0.68) for eGFR decline, ESKD, or renal death | 0.69 (0.53-0.88) | 29.5% vs. 33.9% for total serious adverse events |
| SCORED ³⁵⁾ (CKD with decreased eGFR, T2D) | Sotagliflozin vs. placebo | eGFR 25-60 | 0.71 (0.46-1.08) for eGFR decline or ESKD | 0.99 (0.83-1.18) | 23.4% vs. 25.2% for total serious adverse events |

*Differences were statistically significant (p<0.05) for the presented other outcomes.

ACR, urine albumin-to-creatinine ratio (g/g); CI, confidence interval; DKD, diabetic kidney disease; eGFR, estimated glomerular filtration rate (mL/min/1.73 m²); ESKD, end-stage kidney disease; HR, hazard ratio; s-Cr, serum creatinine; s-K⁺, serum potassium; T2D, type 2 diabetes.

Sodium-glucose cotransporter-2 inhibitors (SGLT2i), which reduce both BP and blood glucose levels²⁵⁻²⁷⁾, have shown a consistent benefit in reducing renal and cardiovascular events in clinical trials²⁵⁻³⁰⁾. Particularly, dapagliflozin, an SGLT2i, has been shown to reduce all-cause mortality as well as the renal and cardiovascular risk in patients with CKD and severe albuminuria, irrespective of the presence

of diabetes³¹⁾. In meta-analyses, the relative risk reduction with SGLT2i was similar among patients with different baseline conditions^{32,33)}. Accordingly, the absolute benefits may be greater in patients at a higher risk of developing renal and cardiovascular events. In a 2021 meta-analysis, the absolute risk reduction in cardiovascular events was greater in patients with a lower eGFR and more severe albuminuria,

and the absolute renoprotective benefit was greater in patients with more severe albuminuria³⁴. In addition, SGLT2i consistently reduced the rate of total serious adverse events in patients with CKD (Table 4)^{29,31,35}. These indicate that the benefits of SGLT2i treatment outweigh the harms in patients with CKD. Furthermore, the net benefit of SGLT2i is greater in patients with CKD, particularly in those with severe albuminuria, than in those without CKD.

Target BP in CKD

Several randomized trials that compared intensive BP control with conventional control in CKD failed to show a benefit in slowing the progression of kidney disease during the trial period³⁶⁻³⁸. However, in extended follow-up studies of the trials, intensive BP control targeting a mean arterial pressure of ≤ 92 mmHg (e.g., BP $\leq 125/75$ mmHg), compared with usual control targeting a mean arterial pressure of ≤ 107 mmHg (e.g., BP $\leq 140/90$ mmHg), was associated with a reduced risk of end-stage kidney disease among patients with CKD and severe proteinuria (Table 1)³⁻⁶.

Meanwhile, in SPRINT, in nondiabetic patients at a high risk of cardiovascular events, targeting an SBP of <120 mmHg, compared with <140 mmHg, resulted in lower rates of cardiovascular events and all-cause mortality⁷. Of the 9,361 SPRINT participants, 2,646 (28%) had an eGFR of <60 mL/min/1.73 m² and 1,723 (19%) had a urine ACR of ≥ 30 mg/g. In subgroup analyses, intensive BP lowering was associated with similar risk reductions in cardiovascular events and all-cause mortality among patients with and without an eGFR of <60 mL/min/1.73 m²^{7,39}. Likewise, the risk reductions in the outcomes of interest with intensive therapy did not clearly differ between patients with and without a urine ACR of ≥ 30 mg/g⁴⁰.

Comparison between SPRINT and Other Studies

In contrast to SPRINT, the Action to Control Cardiovascular Risk in Diabetes trial showed that targeting an SBP of <120 versus <140 mmHg did not reduce all-cause or cardiovascular mortality among patients with type 2 diabetes and a mean eGFR of 92 mL/min/1.73 m²⁴¹. Similarly, in the Heart Outcomes Prevention Evaluation-3 trial, treatment with candesartan

and hydrochlorothiazide did not reduce the risk of composite cardiovascular events among patients with a baseline SBP of ≤ 143.5 mmHg who did not have cardiovascular disease or advanced CKD⁴². Meanwhile, in SPRINT, the relative risk reduction in cardiovascular events with intensive therapy tended to be greater among patients with a baseline SBP of ≤ 132 mmHg than among patients with a baseline SBP of 132 to <145 mmHg and those with an SBP of ≥ 145 mmHg⁷. This is contradictory to the observations that the benefit in terms of BP reduction was greater among patients with higher baseline BP⁴³ and the relative risk reduction was proportional to the magnitude of BP lowering^{44,45}. Assuming that the optimal on-treatment SBP is ≤ 120 mmHg, lowering SBP from a level of 132 to 144 mmHg to a level of ≤ 120 mmHg, compared with lowering SBP from a level of ≤ 132 mmHg, is expected to result in greater risk reduction. Given these discrepancies between SPRINT and other studies, the results of SPRINT should be interpreted with caution.

In addition, randomized trials include participants who are not necessarily representative of a real-world population^{46,47}. Moreover, subgroup analyses are observational in nature and can be influenced by confounders. SPRINT included older adults with a high cardiovascular risk and excluded patients with diabetes or proteinuria (urine protein >1.0 g/day). In addition, there were imbalanced confounders between subgroups with and without CKD (e.g., older adults with CKD). In contrast to the subgroup analyses of SPRINT, a population-based cohort study in Korea matched CKD and non-CKD populations using propensity scores to address potential confounding by imbalanced covariates between the populations. The cohort study revealed that intensively lowered SBP (115 to <125 mmHg), compared with conventionally controlled SBP (135 to <145 mmHg), was associated with a reduced risk of a composite of end-stage kidney disease, cardiovascular events, and all-cause mortality in patients with CKD, particularly in those with albuminuria, but not in their propensity score-matched counterparts without CKD⁴⁸. Considering this finding together with the extended follow-up results of CKD trials (renoprotective effects of intensive BP control were observed in patients with, but not in those without, proteinuria), albuminuria may be an effect modifier of optimal BP, in contrast to the subgroup analysis findings of SPRINT.




Fig. 1. Comparison of all-cause mortality between the lower BP target and standard BP target groups, according to kidney disease status.

A meta-analysis was conducted using the random-effects model. The primary analysis included trials that compared lower BP targets with standard targets (SBP ≤ 140 mmHg, diastolic BP ≤ 90 mmHg, or mean arterial pressure ≤ 107 mmHg) **A**, and secondary analyses were restricted to trials facilitating within-trial comparisons but extended to a trial with a placebo control achieving an SBP of 140 mmHg **B**. The subset “ACCORD-subset- A2,3 and G3” had a urine ACR of ≥ 0.03 in most participants and an eGFR of 30 to < 60 mL/min/1.73 m² in 23.2% of participants. A1, urine ACR < 30 mg/g; A2, urine ACR 30 to 300 mg/g; A3, urine ACR > 300 mg/g; ACR, albumin-to-creatinine ratio; BP, blood pressure; CI, confidence interval; eGFR, estimated glomerular filtration rate; G1, eGFR ≥ 90 mL/min/1.73 m²; G2, eGFR 60 to < 90 mL/min/1.73 m²; G3, eGFR 30 to < 60 mL/min/1.73 m²; G4, eGFR 15 to < 30 mL/min/1.73 m²; SBP, systolic blood pressure.

All-cause Mortality

All-cause mortality can reflect the net benefit of BP reduction. A meta-analysis of clinical trials showed that treatment to a target SBP of < 150 mmHg reduced all-cause mortality (risk ratio [RR] 0.90, 95% confidence interval [CI]

0.83 to 0.98) in adults aged ≥ 60 years with a baseline SBP of ≥ 160 mmHg⁴⁹). Similarly, a 2019 Cochrane review that compared active treatment with no treatment or placebo reported that antihypertensive drug therapy reduced the rates of both all-cause mortality (RR 0.91, 95% CI 0.85 to 0.97) and cardiovascular events (RR 0.72, 95% CI 0.68 to 0.77) in adults aged ≥ 60 years with a baseline BP of

>140/90 mmHg⁵⁰). However, in a 2020 Cochrane review that compared lower BP targets ($\leq 135/85$ mmHg) with standard targets ($\leq 140/90$ mmHg), all-cause mortality did not differ between the lower and standard target groups (RR 0.95, 95% CI 0.86 to 1.05) in the general population of adults with hypertension⁵¹.

In CKD populations, a recent meta-analysis showed that more intensive BP control, compared with less intensive control, was associated with a reduced risk of all-cause mortality (RR 0.86, 95% CI 0.76 to 0.97) in patients with an eGFR of <60 mL/min/1.73 m²⁵². However, the analysis included trials with a control group receiving placebo or no treatment or those targeting a BP higher than standard targets. To compare lower BP targets with standard targets, I conducted a meta-analysis including trials with available data on baseline kidney function that compared lower BP targets with a control targeting an SBP of ≤ 140 mmHg, a diastolic BP of ≤ 90 mmHg, or a mean arterial pressure of ≤ 107 mmHg. I meta-analyzed a total of 9 randomized studies comprising 3 trials in patients with nondiabetic CKD^{36-38,53}, 3 trials in patients with type 2 diabetes^{52,54,55}, a trial in adults with diastolic hypertension^{52,56}, a trial in patients without diabetes with a high cardiovascular risk^{7,40}, and a trial in patients with lacunar stroke⁵⁷, which met the inclusion criteria and had a sample size of ≥ 50 in each target group. I found that more intensive BP control was associated with a lower risk of all-cause mortality in patients with albuminuria (RR 0.76, 95% CI 0.61 to 0.95) but not in those without CKD (Fig. 1A). In patients with an eGFR of ≤ 60 mL/min/1.73 m², intensive versus standard BP control was associated with a reduced risk of all-cause mortality (RR 0.83, 95% CI 0.75 to 0.91) when the extended follow-up data of 2 CKD trials^{5,6} were included, but not when only trial-phase data were analyzed (Fig. 1A). Even when the analysis was restricted to trials facilitating within-trial comparisons but extended to a trial with a placebo control achieving an SBP of 140 mmHg⁵⁸, the results were similar to those of the primary analysis (Fig. 1B). Moreover, in a population-based study in Korea, the optimal on-treatment BP with the best survival outcome was left-shifted in patients with CKD and albuminuria relative to that in patients without CKD⁴⁸. These findings support the assertion that treatment targets lower than standard targets are needed

for the management of BP in patients with CKD.

Adverse Effects

Antihypertensive treatment increases the rate of adverse events and symptoms, including hypotension, dizziness, electrolyte disturbance, arrhythmia, erectile dysfunction, headache, edema, and cough⁴⁹. To achieve lower BP targets, increased numbers and higher doses of antihypertensive drugs are needed. The use of more drugs and higher drug doses can increase the treatment-related adverse effects and the burden on patients. Studies have shown that more intensive BP control increased withdrawals because of adverse events^{49,59,60}. This could lead to higher rates of permanent treatment discontinuation. Treatment to lower BP targets also increased the rate of syncope and other adverse events⁴⁹. In SPRINT, targeting an SBP of <120 mmHg versus <140 mmHg increased the rate of hospitalization or emergency department visits for hypotension, syncope, electrolyte disturbance, and acute kidney injury⁷.

In the 2020 Cochrane review that compared lower BP targets ($\leq 135/85$ mmHg) with standard targets ($\leq 140/90$ mmHg), although the cardiovascular event rates were reduced (RR 0.84, 95% CI 0.73 to 0.96 for myocardial infarction; RR 0.88, 95% CI 0.77 to 1.01 for stroke; and RR 0.75, 95% CI 0.60 to 0.92 for heart failure) in the lower (vs. higher) target group, the rate of other serious adverse events was increased in the lower target group (RR 1.44, 95% CI 1.32 to 1.59) and the rate of total serious adverse events did not differ between the target groups (RR 1.04, 95% CI 0.99 to 1.08)⁵¹. The authors concluded that the benefit of lower BP targets, compared with standard targets, did not outweigh the harms in the general population of persons with high BP.

CONCLUSIONS

Drugs such as SGLT2i, finerenone, and ARNi, in addition to RASi, have shown a benefit in preventing CKD progression in randomized controlled trials. In particular, SGLT2i consistently improved the renal and cardiac outcomes and reduced the rate of total serious adverse events in patients with CKD, and dapagliflozin, an SGLT2i, even reduced all-cause

mortality in patients with severe albuminuria. With respect to the target BP, the KDIGO 2021 BP guideline suggests treating adults with CKD to a target SBP of <120 mmHg based on standardized office BP measurement. This SBP target is largely based on the finding of SPRINT that targeting an SBP of <120 mmHg versus <140 mmHg has both a survival benefit and cardioprotective effects, irrespective of the patient's baseline kidney function. However, previous studies have indicated that albuminuria is an effect modifier of BP targets for renal and mortality outcomes and suggested that a decreased eGFR may be an effect modifier of BP for mortality. Meanwhile, a Cochrane review reported that targeting a BP lower than the standard target ($\leq 140/90$ mmHg) does not further reduce the rate of total serious adverse events despite having a small additional benefit in reducing cardiovascular events. Lowering SBP to <120 mmHg may increase the risk of other serious adverse events beyond the benefit in cardiovascular risk reduction, and a standardized BP measurement increases the burden on patients and resources. Thus, targeting a BP of <130/80 mmHg using an appropriate office BP measurement can be considered an option in patients with CKD. The presence and degree of albuminuria would need to be additionally considered to determine individualized BP targets. Clinical trials evaluating home or ambulatory BP as a treatment target are warranted, and more studies are needed to confirm the role of albuminuria as an effect modifier of BP for overall outcomes.

Acknowledgment

I specially thank Prof. Gheun-Ho Kim at Hanyang University College of Medicine for his helpful comments during manuscript drafting.

Conflict of Interest

The authors have no conflicts of interest to declare.

ORCID: Hae Hyuk Jung 0000-0002-3202-6307

REFERENCES

- Cheung AK, Chang TI, Cushman WC, et al. Executive summary of the KDIGO 2021 Clinical Practice Guideline for the Management of Blood Pressure in Chronic Kidney Disease. *Kidney Int.* 2021;99:559-569.
- Wheeler DC, Becker GJ. Summary of KDIGO guideline. What do we really know about management of blood pressure in patients with chronic kidney disease? *Kidney Int.* 2013; 83:377-383.
- Sarnak MJ, Greene T, Wang X, et al. The effect of a lower target blood pressure on the progression of kidney disease: long-term follow-up of the modification of diet in renal disease study. *Ann Intern Med.* 2005;142:342-351.
- Appel LJ, Wright JT, Greene T, et al. Intensive blood-pressure control in hypertensive chronic kidney disease. *N Engl J Med.* 2010;363:918-929.
- Ku E, Glidden DV, Johansen KL, et al. Association between strict blood pressure control during chronic kidney disease and lower mortality after onset of end-stage renal disease. *Kidney Int.* 2015;87:1055-1060.
- Ku E, Gassman J, Appel LJ, et al. BP Control and Long-Term Risk of ESRD and Mortality. *J Am Soc Nephrol JASN.* 2017; 28:671-677.
- SPRINT Research Group, Wright JT Jr, Williamson JD, et al. A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med.* 2015;373:2103-2116.
- Maschio G, Alberti D, Janin G, et al. Effect of the angiotensin-converting-enzyme inhibitor benazepril on the progression of chronic renal insufficiency. The Angiotensin-Converting-Enzyme Inhibition in Progressive Renal Insufficiency Study Group. *N Engl J Med.* 1996;334:939-945.
- Ruggenenti P, Perna A, Gherardi G, et al. Renoprotective properties of ACE-inhibition in non-diabetic nephropathies with non-nephrotic proteinuria. *Lancet Lond Engl.* 1999; 354:359-364.
- Randomised placebo-controlled trial of effect of ramipril on decline in glomerular filtration rate and risk of terminal renal failure in proteinuric, non-diabetic nephropathy. The GISEN Group (Gruppo Italiano di Studi Epidemiologici in Nefrologia). *Lancet Lond Engl.* 1997;349:1857-1863.
- Ruggenenti P, Perna A, Remuzzi G. ACE inhibitors to prevent end-stage renal disease: when to start and why possibly never to stop: a post hoc analysis of the REIN trial results. Ramipril Efficacy in Nephropathy. *J Am Soc Nephrol JASN.* 2001;12:2832-2837.
- Hou FF, Zhang X, Zhang GH, et al. Efficacy and safety of benazepril for advanced chronic renal insufficiency. *N Engl J Med.* 2006;354:131-140.
- Lewis EJ, Hunsicker LG, Bain RP, Rohde RD. The effect of angiotensin-converting-enzyme inhibition on diabetic nephropathy. The Collaborative Study Group. *N Engl J Med.* 1993;

- 329:1456-1462.
14. Brenner BM, Cooper ME, de Zeeuw D, et al. Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med.* 2001;345:861-869.
 15. Lewis EJ, Hunsicker LG, Clarke WR, et al. Renoprotective effect of the angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Engl J Med.* 2001;345:851-860.
 16. Agodoa LY, Appel L, Bakris GL, et al. Effect of ramipril vs amlodipine on renal outcomes in hypertensive nephrosclerosis: a randomized controlled trial. *JAMA.* 2001;285:2719-2728.
 17. Telmisartan, Ramipril, or Both in Patients at High Risk for Vascular Events. *N Engl J Med.* 2008;358:1547-1559.
 18. Fried LF, Emanuele N, Zhang JH, et al. Combined Angiotensin Inhibition for the Treatment of Diabetic Nephropathy. *N Engl J Med.* 2013;369:1892-1903.
 19. Parving H-H, Brenner BM, McMurray JJV, et al. Cardiorenal End Points in a Trial of Aliskiren for Type 2 Diabetes. *N Engl J Med.* 2012;367:2204-2213.
 20. McMurray JJV, Krum H, Abraham WT, et al. Aliskiren, Enalapril, or Aliskiren and Enalapril in Heart Failure. *N Engl J Med.* 2016;374:1521-1532.
 21. Bakris GL, Agarwal R, Anker SD, et al. Effect of Finerenone on Chronic Kidney Disease Outcomes in Type 2 Diabetes. *N Engl J Med.* 2020;383:2219-2229.
 22. Pitt B, Filippatos G, Agarwal R, et al. Cardiovascular Events with Finerenone in Kidney Disease and Type 2 Diabetes. *N Engl J Med.* 2021;0:null.
 23. Solomon SD, McMurray JJV, Anand IS, et al. Angiotensin-Nephrilysin Inhibition in Heart Failure with Preserved Ejection Fraction. *N Engl J Med.* Published online September 1, 2019.
 24. McMurray JJV, Packer M, Desai AS, et al. Angiotensin-Nephrilysin Inhibition versus Enalapril in Heart Failure. *N Engl J Med.* 2014;371:993-1004.
 25. Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med.* 2015;373:2117-2128.
 26. Neal B, Perkovic V, Mahaffey KW, et al. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med.* 2017;377:644-657.
 27. Wiviott SD, Raz I, Bonaca MP, et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med.* Published online November 10, 2018.
 28. Wanner C, Inzucchi SE, Lachin JM, et al. Empagliflozin and Progression of Kidney Disease in Type 2 Diabetes. *N Engl J Med.* 2016;375:323-334.
 29. Perkovic V, Jardine MJ, Neal B, et al. Canagliflozin and Renal Outcomes in Type 2 Diabetes and Nephropathy. *N Engl J Med.* Published online April 14, 2019.
 30. McMurray JJV, Solomon SD, Inzucchi SE, et al. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med.* 2019;381:1995-2008.
 31. Heerspink HJL, Stefánsson BV, Correa-Rotter R, et al. Dapagliflozin in Patients with Chronic Kidney Disease. *N Engl J Med.* 2020;383:1436-1446.
 32. Zelniker TA, Wiviott SD, Raz I, et al. SGLT2 inhibitors for primary and secondary prevention of cardiovascular and renal outcomes in type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet Lond Engl.* 2019;393:31-39.
 33. Neuen BL, Young T, Heerspink HJL, et al. SGLT2 inhibitors for the prevention of kidney failure in patients with type 2 diabetes: a systematic review and meta-analysis. *Lancet Diabetes Endocrinol.* 2019;7:845-854.
 34. Chun KJ, Jung HH. SGLT2 Inhibitors and Kidney and Cardiac Outcomes According to Estimated GFR and Albuminuria Levels: A Meta-analysis of Randomized Controlled Trials. *Kidney Med.* 2021;0.
 35. Bhatt DL, Szarek M, Pitt B, et al. Sotagliflozin in Patients with Diabetes and Chronic Kidney Disease. *N Engl J Med.* 2020;0:null.
 36. Klahr S, Levey AS, Beck GJ, et al. The effects of dietary protein restriction and blood-pressure control on the progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. *N Engl J Med.* 1994;330:877-884.
 37. Jackson T, Wright J, Bakris G, Greene T, et al. Effect of Blood Pressure Lowering and Antihypertensive Drug Class on Progression of Hypertensive Kidney Disease: Results From the AASK Trial. *JAMA.* 2002;288:2421-2431.
 38. Ruggenenti P, Perna A, Loriga G, et al. Blood-pressure control for renoprotection in patients with non-diabetic chronic renal disease (REIN-2): multicentre, randomised controlled trial. *Lancet Lond Engl.* 2005;365:939-946.
 39. Cheung AK, Rahman M, Reboussin DM, et al. Effects of Intensive BP Control in CKD. *J Am Soc Nephrol JASN.* 2017; 28:2812-2823.
 40. Chang AR, Kramer H, Wei G, et al. Effects of Intensive Blood Pressure Control in Patients with and without Albuminuria: Post Hoc Analyses from SPRINT. *Clin J Am Soc Nephrol.* 2020;15:1121-1128.
 41. ACCORD Study Group, Cushman WC, Evans GW, et al. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med.* 2010;362:1575-1585.

42. Lonn EM, Bosch J, López-Jaramillo P, et al. Blood-Pressure Lowering in Intermediate-Risk Persons without Cardiovascular Disease. *N Engl J Med*. 2016;374:2009-2020.
43. Brunström M, Carlberg B. Association of Blood Pressure Lowering With Mortality and Cardiovascular Disease Across Blood Pressure Levels: A Systematic Review and Meta-analysis. *JAMA Intern Med*. 2018;178:28-36.
44. Ettehad D, Emdin CA, Kiran A, et al. Blood pressure lowering for prevention of cardiovascular disease and death: a systematic review and meta-analysis. *Lancet*. 2016;387:957-967.
45. Bangalore S, Toklu B, Gianos E, et al. Optimal Systolic Blood Pressure Target After SPRINT: Insights from a Network Meta-Analysis of Randomized Trials. *Am J Med*. 2017;130:707-719.e8.
46. Stuart EA, Bradshaw CP, Leaf PJ. Assessing the Generalizability of Randomized Trial Results to Target Populations. *Prev Sci Off J Soc Prev Res*. 2015;16:475-485.
47. Kennedy-Martin T, Curtis S, Faries D, Robinson S, Johnston J. A literature review on the representativeness of randomized controlled trial samples and implications for the external validity of trial results. *Trials*. 2015;16:495.
48. Jung HH. On-treatment blood pressure and long-term outcomes in chronic kidney disease. *Nephrol Dial Transplant*. 2021.
49. Qaseem A, Wilt TJ, Rich R, et al. Pharmacologic Treatment of Hypertension in Adults Aged 60 Years or Older to Higher Versus Lower Blood Pressure Targets: A Clinical Practice Guideline From the American College of Physicians and the American Academy of Family Physicians. *Ann Intern Med*. 2017;166:430-437.
50. Musini VM, Tejani AM, Bassett K, Puil L, Wright JM. Pharmacotherapy for hypertension in adults 60 years or older. *Cochrane Database Syst Rev*. 2019.
51. Arguedas JA, Leiva V, Wright JM. Blood pressure targets in adults with hypertension. *Cochrane Database Syst Rev*. 2020.
52. Malhotra R, Nguyen HA, Benavente O, et al. Association Between More Intensive vs Less Intensive Blood Pressure Lowering and Risk of Mortality in Chronic Kidney Disease Stages 3 to 5: A Systematic Review and Meta-analysis. *JAMA Intern Med*. 2017;177:1498-1505.
53. Aggarwal Rahul, Petrie Benjamin, Bala Wasif, Chiu Nicholas. Mortality Outcomes With Intensive Blood Pressure Targets in Chronic Kidney Disease Patients. *Hypertension*. 2019;73:1275-1282.
54. Schrier RW, Estacio RO, Mehler PS, Hiatt WR. Appropriate blood pressure control in hypertensive and normotensive type 2 diabetes mellitus: a summary of the ABCD trial. *Nat Clin Pract Nephrol*. 2007;3:428-438.
55. Papademetriou V, Zaheer M, Doumas M, et al. Cardiovascular Outcomes in Action to Control Cardiovascular Risk in Diabetes: Impact of Blood Pressure Level and Presence of Kidney Disease. *Am J Nephrol*. 2016;43:271-280.
56. Hansson L, Zanchetti A, Carruthers SG, et al. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. *The Lancet*. 1998;351:1755-1762.
57. Blood-pressure targets in patients with recent lacunar stroke: the SPS3 randomised trial. *The Lancet*. 2013;382:507-515.
58. Heerspink HJL, Ninomiya T, Perkovic V, et al. Effects of a fixed combination of perindopril and indapamide in patients with type 2 diabetes and chronic kidney disease. *Eur Heart J*. 2010;31:2888-2896.
59. Thomopoulos C, Parati G, Zanchetti A. Effects of blood pressure lowering treatment in hypertension: 8. Outcome reductions vs. discontinuations because of adverse drug events—meta-analyses of randomized trials. *J Hypertens*. 2016;34:1451-1463.
60. Saiz LC, Gorricho J, Garjón J, Celaya MC, Erviti J, Leache L. Blood pressure targets for the treatment of people with hypertension and cardiovascular disease. *Cochrane Database Syst Rev*. 2020.